

1
2
3
4 1
5
6 2 **Two-dimensional proteome reference map of *Vibrio tapetis*, the etiological**
7
8 **agent of brown ring disease in clams**
9 3
10
11 4

12
13
14 5 José Bermúdez-Crespo¹, Sabela Balboa², Jana Alonso³ and Jesús L. Romalde² *

15 6
16
17
18 7 ¹Departamento de Genética, and ²Departamento de Microbiología y Parasitología. CIBUS-
19 8 Facultad de Biología, Universidad de Santiago de Compostela. Campus Vida s/n. 15782,
20 9 Santiago de Compostela. Spain.

21 10 ³Servicio de Proteómica, Fundación IDICHUS, Hospital Clínico Universitario. Travesía da
22 11 Choupana s/n. 15706. Santiago de Compostela, Spain.
23
24

25 12
26 13 **Running title:** Description of the reference map of *Vibrio tapetis*.
27
28
29
30
31

32 14
33
34
35
36 15 Submitted to: **Journal of Applied Microbiology**, February 2012

37
38 16 Revised Ms. JAM-2011-2115
39
40
41
42

43 18 *Corresponding author:
44

45 19 Departamento de Microbiología y Parasitología. CIBUS-Facultad de Biología, Universidad de
46 20 Santiago de Compostela. Campus Vida s/n. 15782, Santiago de Compostela. Spain.
47
48

49 21 Telf.: (+34) 881816908

50 22 Fax: (+34) 881816966

51 23 E-mail: jesus.romalde@usc.es
52
53
54
55
56
57
58
59
60

1
2
3
4 255
6 **ABSTRACT**7
8 27

9
10 28 *Vibrio tapetis* is the etiological agent of brown ring disease (BRD) in clams, one of the most
11
12 29 threatening diseases affecting this commercially important bivalve. In this study we have
13
14 30 constructed a proteome reference map of the *V. tapetis* type strain CECT 4600^T. Eighty-two
15
16 31 proteins, consistently present in all gels, were identified by mass spectrometry or by *de novo*
17
18 32 sequencing. The majority of the proteins identified (66%) belonged to four COG categories:
19
20 33 “Carbohydrate transport and metabolism”, “Post-translational modification, protein turnover
21
22 34 and chaperones”, “Energy production”, and “Amino acid transport and metabolism”.
23
24 35 Glyceraldehyde-3-phosphate dehydrogenase, enolase, fructose-bisphosphate aldolase,
25
26 36 phosphoglycerate kinase, molecular chaperones DnaK and GroEL, alkyl hydroperoxide
27
28 37 reductase, peptidyl-prolyl cis-trans isomerase B and factor Tu, were identified among the 20
29
30 38 most abundant proteins. A comparison of this reference map with that obtained for the *V.*
31
32 39 *tapetis* strain GR0202RD, with different origin and pathophysiological characteristics, was
33
34 40 performed. The results obtained here can be considered as a first step to gather valuable
35
36 41 information on protein expression, related not only to diverse cellular functions and regulation
37
38 42 but also to pathogenesis and bacterium-host interactions in the disease process.
39
40 43
41 44
42 45
43 46
44 47
45 48

46 49 **Key words:** *Vibrio tapetis*, proteome, reference map, MALDI-TOFF, *de novo* sequencing
47 50
48 51
49 52
50 53
51 54
52 55
53 56
54 57
55 58
56 59
57 60

48 INTRODUCTION

49 Brown Ring Disease (BRD), caused by *Vibrio tapetis* (Borrego *et al.* 1996), is an epizootic
50 infection described in adult clams. The main sign characterizing the disease is a brown
51 conchiolin deposit on the inner surface of the valves, typically located between the pallial line
52 and the edge of the shell. This organic deposit perturbs the calcification process (Paillard
53 2004; Paillard *et al.* 1994; Trinkler *et al.* 2010; Trinkler *et al.* 2011) causing severe
54 deformations of the clam's shell and subsequently a systemic infection that provokes the
55 death of the animal. Three groups has been described within *V. tapetis*, on the basis of the
56 phenotypic, antigenic, and genotypic characteristics of the isolates, that correlate with the host
57 origin (Castro *et al.* 1996; Castro *et al.* 1997; Le Chevalier *et al.* 2003; Rodríguez *et al.* 2006;
58 Romalde *et al.* 2002).

59 The pathogenicity process is not well established, although it has described that not all *V.*
60 *tapetis* strains have the same capacity of producing BRD signs and/or mortalities (Novoa *et*
61 *al.* 1998; Choquet 2004; Paillard 2004). In general, *V. tapetis* shows adhesion capability
62 toward the clam hemocytes and mantle cells (López-Cortés *et al.* 1999) and an *in vitro*
63 experiment has demonstrated that the cytotoxic effect on clam hemocytes results in cell
64 rounding and the loss of filopods (Lane and Birbeck 2000; Choquet *et al.* 2003).

65 Some adherence factors, such as pili, have been identified and it is known that they allow the
66 colonization and proliferation of this pathogen in the peripheral compartment (Paillard and
67 Maes 1995; Paillard 2004). Hemolysin and exo-enzymes, such as esterase or chymotrypsin,
68 have been detected as extracellular products with a role in pathogenesis (Borrego *et al.* 1996;
69 Allam *et al.* 2001; Allam *et al.* 2002; Allam and Ford 2006). More recently, it was
70 demonstrated that DjlA (DnaJ-like protein A), an inner membrane-anchored bacterial
71 cochaperone, is involved in *V. tapetis* cytotoxicity (Lakhal *et al.* 2008), since a DjlA defective

1
2
3
4 72 mutant showed marked reduction in cytotoxicity activity, compared with the wild type strain,
5
6 73 against *R. philippinarum* hemocytes *in vitro*.
7

8
9 74 The description of a proteomic reference map can provide valuable data for the protein
10
11 75 expression, related to diverse cellular functions and regulation (Phillips and Bogyo 2005; Cash
12
13 76 2009; Cash and Argo 2009), the pathogenesis pathways and the bacterium-host interactions in
14
15 77 the disease process (Osman *et al.* 2009; Bumann 2010), as well as the response of a given
16
17 78 organism to various external stimuli (Jones *et al.* 2008; Poobalane *et al.* 2008; Roncada *et al.*
18
19 79 2009; Sharma *et al.* 2010). In the last years, the proteomes of some vibrios have been
20
21 80 analyzed and several proteins possibly linked to pathogenesis have been characterized
22
23 81 (Coelho *et al.* 2004; Kan *et al.* 2004; Lee *et al.* 2006). The knowledge of such reference map
24
25 82 is even more important in the case of pathogens as *V. tapetis*, in which the complete genome
26
27 83 was not yet published. This study represents the first description of the reference map of *V.*
28
29 84 *tapetis*.
30
31
32
33
34

35 86 MATERIAL AND METHODS

37 87 Bacterial strains

38
39 88 The type strain CECT 4600^T of *Vibrio tapetis*, obtained from Manila clam (*Ruditapes*
40
41 89 *philippinarum*), as well as strain GR0202RD isolated from carpet-shell clam (*R. decussatus*)
42
43 90 in Spain, showing different physiological and genetic characteristics and virulence capacity,
44
45 91 were used in this study (Borrego *et al.* 1996; Novoa *et al.* 1998; Rodríguez *et al.* 2006). The
46
47 92 bacteria were routinely grown on Marine Agar (MA) (Pronadisa, Spain) under aerobic
48
49 93 conditions at 15°C for 72h. Stock cultures were maintained frozen at -80°C in Marine Broth
50
51 94 (MB) (Pronadisa, Spain) supplemented with 15% glycerol (v/v).
52
53
54
55
56
57
58
59
60

96 **2DE-PAGE**

97 *Growth conditions*

98 Bacterial inocula were prepared resuspending each strain to achieve 10^9 cells per milliliter
99 adjusting the bacterial suspension to an OD=1 (508 nm) using a Lambda2 UV/VIS
100 Spectrophotometer (Perkin Elmer).

101 One flask per strain containing 1 liter of sterile Marine Broth (MB, Pronadisa, Spain) was
102 inoculated with 250 μ l of a 10^8 cells/ml tube achieving a final concentration of 10^5 cells/ml
103 and aerobically incubated in a Innova 4340 rotary shaker (70 rpm) (New Brunswick
104 Scientific, USA) at 15°C for 72 hours.

105 *Sample preparation for 2DE-PAGE*

106 Bacteria were harvested and washed with Tris-buffered sucrose (10 mmol Tris, 250 mmol
107 sucrose pH 7) and lyophilised. Proteins were extracted by suspending 40 mg of lyophilised
108 bacteria in 1 mL standard lysis buffer (7M urea, 2M thiourea, 4% CHAPS (3-[(3-
109 cholamidopropyl) dimethylammonio]-1-propanesulfonate) and 65 mM DTT (Dithiothreitol)
110 during 3 hours at 27 °C. In the middle of digestion, samples were sonicated with three cycles
111 of 10 pulses with 15 seconds of separation between each cycle. Next, samples were
112 centrifuged at 14000 rpm for 30 min. The resulting supernatants were collected and subjected
113 to protein precipitation using the Clean-up kit (GE Healthcare, USA). After pellet re-
114 suspension in other 1 mL of lysis buffer, protein concentration were measured with CB-X
115 protein assay kit (Gbiosciences, USA). Finally, samples were stored at -80 °C prior to use.

116 *First dimension*

117 Isoelectrofocusing (IEF) was performed using a Protean IEF cell (Bio-Rad, USA) and 24 cm
118 pH 4-7 IPG strips (GE Healthcare, Sweden). Each 60 μ l sample (containing approximately
119 400 μ g of protein) was mixed with 390 μ l of rehydration buffer (7M urea, 2M thiourea, 4%

1
2
3
4 120 CHAPS, 0.6 % DTT, 1% IPG buffer 4-7 and bromophenol blue traces) and loaded in each
5
6 121 strip by in-gel-rehydration (18). IEF was carried out at 20 °C and according to the following
7
8 122 steps: active rehydration (50V) for 12h, 250 V for 30 min, 500 V for 1h, 1000 V for 1h, 4000
9
10 123 V for 2h, 8000 V for 2 h and 10000 V to achieve 65 kVh. After IEF, strips were stored at -20
11
12 124 °C until used in second dimension.

125 *Second dimension*

126 Prior to run the second dimension, strips were equilibrated at room temperature in
127 equilibration solution (6M urea, 50mM Tris-HCl pH 8.8, 30% glycerol, 2% SDS) with
128 addition of 1% DTT for 15 min, followed by the same solution with addition of 2.5% of
129 iodoacetamide other 15 min. Strips were fixed with sealed solution (25mM Tris, 192 mM
130 glycine, 0.1% SDS, 0.5% agarose, 0.01% bromophenol blue) on top of a 21 x 26 cm 12,5%
131 polyacrilamide gel. Second dimension was performed according to Laemmli (Laemmli 1970),
132 in a EttanDalt-Six (GE Healthcare PAIS) at 30° C overnight (5mA/gel for 1 h, 10mA/per 1 h,
133 followed by 16 mA/gel until bromophenol blue dye reached the bottom of the gel).

134 Two independent cultures as well as three replicates gel per culture were performed to assess
135 the reproducibility of the experiment.

136 *Protein visualization and analysis*

137 Gels were stained with Coomassie Brilliant Blue (CBB). CBB staining was carried out
138 according to Neuhoff's protocol with minor modifications (Neuhoff *et al.* 1988). Gels were
139 fixed in 50% ethanol, 3% phosphoric acid for 3 h and then washed in H₂O three times for 20
140 min each. The gels were pre-incubated in 34% methanol, 17% sulphate ammonium, 3%
141 phosphoric acid for 1 h. Next, 0.35 g/L of Coomassie Blue G-250 (Bio-Rad) was added to this
142 solution. Each gel was stained alone in a plastic box with 330 mL of this solution during four
143 days to reach maximum intensity of staining. Finally, gels were washed in H₂O twice for 25

1
2
3
4 144 min each and scanned in a densitometer at 300 dpi resolution. Digitalised images were saved
5
6 145 in TIFF format and gels were stored in vacuum-sealed plastic bags at 4°C. PDQuest Advance
7
8 146 software version 8.0 (Bio-Rad, USA) were used for spot detection, spot quantitation and
9
10 147 reproducibility study.
11
12
13 148

149 **Mass spectrometric analysis**

150 Protein features chosen for mass spectrometric analysis were excised from the gels and
151 manually digested as previously described (De La Fuente *et al.* 2011). Dried samples were
152 dissolved in 4 µL of 0.5% HCOOH. Equal volumes (0.5 µL) of peptide and matrix solution,
153 consisted of 3 mg CHCA dissolved in 1 mL of 50% acetonitrile in 0.1% TFA, were deposited
154 using the thin layer method, onto a 384 Opti-TOF MALDI plate (Applied Biosystems). Mass
155 spectrometric data were obtained in an automated analysis loop using 4800 MALDI-
156 TOF/TOF analyzer (Applied Biosystems). MS spectra were acquired in reflectron positive-
157 ion mode with a Nd:YAG, 355 nm wavelength laser, averaging 1000 laser shots and at least
158 three trypsin autolysis peaks were used as internal calibration. All MSMS spectra were
159 performed by selection the precursors with a relative resolution of 300 (FWHM) and
160 metastable suppression. Automated analysis of mass data was achieved using the 4000 Series
161 Explorer Software V3.5. MS and MSMS spectra data were combined through the GPS
162 Explorer Software v3.6 using Mascot software v2.1. (Matrix Science) to search against a non-
163 identical protein database (NCBI nr release data 20100526), with 30 ppm precursor tolerance,
164 0.35 Da MSMS fragment tolerance, CAM (carbamylmethyl cystein) as fixed modification,
165 oxidized methionine as variable modification and allowing 1 missed cleavage. All spectra and
166 database results were manually inspected in detail using the previous software. Protein scores
167 greater than 56 were accepted as significant ($p < 0.05$), considering positive the identification

1
2
3
4 168 when protein score CI% (Confidence Interval) was above 98. In case of MSMS spectra, total
5
6 169 ion score CI% was above 95. Spots not identified with Mascot search engine in the automatic
7
8 170 loop were submitted to de novo sequencing. A maximum of 8 peptides were launched
9
10 171 combining all spectra, allowing a mass tolerance of 0.2 Da and PAM30MS as scoring matrix
11
12 172 (4).

173 **RESULTS**

174 Three replicates of the proteome of two independent cultures of *Vibrio tapetis* CECT 4600^T
175 strain were separate on large 2-DE gels in *pI* range of 4-7. Gels within each culture as well as
176 between both cultures showed high reproducibility (Fig. 1). Automatic software analysis was
177 used to determine the spot number in each gel. While total amount of protein loaded on each
178 gel was the same, approximately 400 µg, there was significant variation in spot number
179 between both cultures. Thus, average number (\pm SD) of spot detected was 428 (\pm 7) and 504
180 (\pm 9) for first and second culture, respectively. Most of the proteins detected were localised in
181 the acidic part of the pH range studied (Fig. 1 & 2). Visual inspection also revealed different
182 protein expression according to different molecular mass region. Thus, most abundant
183 proteins were localised, very close, in high molecular weight region, whereas faintest proteins
184 were localised, more expanded, in low molecular weight region (Fig. 1 & 2).

185 From all visualised proteins, 82 proteins that were consistently present in all gels were
186 identified by mass spectrometry (Fig. 2). Sixty-nine of these 82 proteins were satisfactorily
187 identified by MS or MS/MS, representing 61 different unique proteins (Table 1 and
188 Supplementary table1). More than one identification per spot was only obtained for one spot
189 (spot 25) (Table1). Unidentified proteins (13 proteins) were subjected to *de novo* sequencing
190 to obtain information about their identity (Table 2 and Supplementary table 2). Identified
191 proteins were classified according to their biological functions using the NCBI COG database

1
2
3
4 192 (<http://www.ncbi.nlm.nih.gov/COG/old/>). Most of the identified proteins were involved in
5
6 193 “Cellular processes” and “Metabolism” classes (85% of all identified proteins). Proteins
7
8 194 involved in “Information, storage and processing” were also identified and only three proteins
9
10 195 were poorly characterized (Fig. 3). More than a half (66%) of the proteins belonging to the
11
12 196 cellular processes and metabolism classes, could be grouped within four of the 13 possible
13
14 197 COG functional categories: i) “Carbohydrate transport and metabolism” (G), ii) “Post-
15
16 198 translational modification, protein turnover and chaperones” (O), iii) “Energy production”
17
18 199 (C), and iv) “Amino acid transport and metabolism” (E). The rest of the identified proteins
19
20 200 were related with other COG categories, including “Inorganic ion transport and metabolism”
21
22 201 (P), “Cell envelope biogenesis and outer membrane” (M), “Signal transduction mechanism”
23
24 202 (T), “Translation, ribosomal structure and biogenesis” (J), “Transcription” (K), “DNA
25
26 203 replication, recombination and repair” (L), “Lipid metabolism” (I), “Coenzyme metabolism”
27
28 204 (H) and “Nucleotide transport and metabolism” (F) (Fig. 3).

29
30
31
32
33 205 The 20 most abundant proteins are listed in Table 3. Among these proteins, 40% were
34
35 206 involved in “Cellular processes”, 35% in “Metabolism”, 20% in “Information, storage and
36
37 207 processing”, and 5% were not identified. In terms of COG functional categories, categories C
38
39 208 (6 proteins) G (4 proteins), and O (4 proteins) were the most represented.

40
41 209 The most abundant proteins in the COG functional category C (energy production and
42
43 210 conversion) were phosphoenolpyruvate carboxykinase, malate dehydrogenase and ATPase
44
45 211 synthetase enzyme, which was detected in two different spots being one of them the beta
46
47 212 subunit. The alpha and beta subunits of Succinyl-CoA synthetase were also identified. COG
48
49 213 category G (carbohydrate transport and metabolism) was represented by four proteins
50
51 214 involved in the glycolysis/gluconeogenesis pathway: glyceraldehyde-3-phosphate
52
53 215 dehydrogenase, enolase, fructose-bisphosphate aldolase, and phosphoglycerate kinase. In the
54
55
56
57
58
59
60

1
2
3
4 216 COG category O (post-translational modification, protein turnover and chaperones) were
5
6 217 included the molecular chaperones Dnak and GroEL (hsp60 family), alkyl hydroperoxide
7
8 218 reductase and peptidyl-prolyl cis-trans isomerase B.

9
10 219 However, the second most abundant protein in the gels belonged to the COG category
11
12 220 “Translation, ribosomal structure and biogenesis” (J). It was identified as the elongation factor
13
14 221 Tu, which is an essential protein for protein synthesis.
15
16

17
18
19 222
20 223 The proteome of *V. tapetis* strain GR0202RD, was also analysed under the same conditions.

21
22 224 Fifty six spots with similar position in both strains were detected. All these proteins were
23
24 225 subjected to MS in order to assess their identity. In addition, another 27 proteins were
25
26 226 subjected to MS to obtain more information about the proteome of strain GR0202RD (Fig. 4).

27
28 227 From all 56 possible similar proteins, 45 (80%) matched in both position and identity, and
29
30 228 only 3 proteins that matched in position had a different identity (Table 4). For the rest of
31
32 229 proteins, this analysis could not be performed since one or both proteins of the pairs were not
33
34 230 identified by MS (Table 4).

35
36
37 231 Of the 20 most abundant proteins in the type strain CECT 4600^T (Table 3), 18 were also
38
39 232 identified in GR0202RD strain (with the exception of spots 27 and 32). On the other hand,
40
41 233 new proteins identified in GR0202RD were classified into COG functional categories.

42
43 234 Although GR0202RD strain did not contribute with any new protein in the most represented
44
45 235 COG category (carbohydrate transport and metabolism), a new COG category, “Cell division
46
47 236 and chromosome partitions” (D), was observed with the identification of the chromosome
48
49 237 segregation ATPase.
50

51
52
53 238
54
55
56
57
58
59
60

1
2
3
4 239 The most abundant protein in each strain (corresponding to spots 37 and 104) could not be
5
6 240 identified by conventional MS or MS/MS. Although in different position in the gels, both
7
8 241 spots were identified as the Porin-like protein H (OmpH) by *de novo* sequencing.
9

10 242

11 12 13 243 **DISCUSSION**

14
15 244 *Vibrio tapetis* is an opportunistic pathogen, which has caused important lost in the clam
16
17 245 culture all over the world (Paillard 2004). The development of BRD is the result of complex
18
19 246 interactions between the pathogen, its host and the environment and, in fact *V. tapetis* has
20
21 247 been proposed as a model to study the host-pathogen interactions (Paillard 2011). Since the
22
23 248 complete genome of this microorganism, although under preparation (C. Paillard, personal
24
25 249 communication), is not yet published, the proteomic approach may help to clarify its
26
27 250 interactions with the clam and the disease processes. In this work, the proteome reference map
28
29 251 of *V. tapetis* is described under laboratory conditions using two strains of the species with
30
31 252 remarkable differences at genetic and pathophysiological levels (Borrego *et al.* 1996; Novoa
32
33 253 *et al.* 1998; Rodríguez *et al.* 2006).

34
35
36
37 254 A high reproducibility was obtained for the protein profiles of the *V. tapetis* strains, the
38
39 255 majority of the spots being detected in the acidic part of the 2D-gels. This finding is in
40
41 256 agreement with results of other authors observing a predominance of proteins with low *pI*
42
43 257 over high *pI* in halophilic bacteria (Kiraga *et al.* 2007).

44
45
46 258 Similar to other bacterial proteomes (Coelho *et al.* 2004; Chung *et al.* 2007), most of the
47
48 259 identified proteins were involved in “Cellular processes” and “Metabolism” classes and in a
49
50 260 lesser extent in “Information, storage and processing” class. The most represented COG
51
52 261 functional categories were “Carbohydrate transport and metabolism” (G), “Post-translational
53
54 262 modification, protein turnover and chaperones” (O), “Energy production” (C) and “Amino
55
56
57
58
59
60

1
2
3
4 263 acid transport and metabolism” (E). These results are logically expected since within these
5
6 264 categories there are very important pathways, which are essential for the cell survival.
7
8
9

10
11 266 The most abundant detected protein was phosphoenolpyruvate carboxykinase which although
12
13 267 is related to COG category “Energy production and conversion”, has also a main role in the
14
15 268 gluconeogenesis, which is related to the COG “Carbohydrate transport and metabolism”
16
17 269 pathway. Phosphoenolpyruvate carboxykinase converts oxalacetate into phosphoenolpyruvate
18
19 270 and carbon dioxide, and play an important role in the reversion of the irreversible reaction of
20
21 271 pyruvate kinase (Hanson and Patel 1994). The second most abundant protein is the elongation
22
23 272 factor Tu, which is an essential protein for protein synthesis. Alkyl hydroperoxide reductase
24
25 273 was the third most abundant protein, but its high expression was unexpected because it takes
26
27 274 part in the oxidative defence system of the cell in situations with high levels of H₂O₂
28
29
30
31 275 (VanBogelen *et al.* 1987).

32
33 276 Malate dehydrogenase was also identified in the 2D gels. This enzyme is involved in the TCA
34
35 277 cycle, but also in the gluconeogenesis since it catalyzes the conversion of malate into
36
37 278 oxalacetate (Gietl 1992). It is noteworthy the finding into the top 20 most abundant proteins
38
39 279 of two enzymes that are linked together. Thus, the product of the reaction catalyzed by one
40
41 280 enzyme serves as substrate for the second enzyme. In this case, oxalacetate, the product of
42
43 281 malate dehydrogenase reaction, serves as substrate for the reaction catalyzed by
44
45 282 phosphoenolpyruvate carboxykinase in the gluconeogenesis pathway.

46
47
48 283 ATPase synthetase enzyme, detected in two different spots, is the major producer of ATP in
49
50 284 bacterial cells. One of these identifications was the beta subunit of ATP synthetase, which is
51
52 285 important as a main component of the second stalk connecting the F1 and F0 sectors, being
53
54 286 essential for normal assembly and function (Dunn *et al.* 2000). The detected alpha and beta
55
56
57
58
59
60

1
2
3
4 287 subunits of Succinyl-CoA synthetase play a role in the formation of succinate and coenzyme-
5
6 288 A from succinyl-CoA in a reversible reaction in the TCA cycle. In addition, this reaction
7
8 289 generates one molecule of GTP, which transfers the phosphate group to ADP to generate one
9
10 290 ATP (32). On the other hand, high expression of peptidyl-prolyl cis-trans isomerase B, Dnak
11
12 291 and GroEL was not surprising since they are essential for protein folding in the cell (Hayano
13
14 292 *et al.* 1991; Gragerov *et al.* 1992).

15
16
17 293 For comparative purposes, the proteome of strain GR0202RD was also characterized. This
18
19 294 strain shows different serological properties than CECT 4600^T strains as evidenced by slide
20
21 295 agglutination and dot-blot assays (Novoa *et al.* 1998; López *et al.* 2010). In addition, they
22
23 296 belong to different genogroups within *V. tapetis* as demonstrated employing several PCR
24
25 297 typing methods and ribotyping (Romalde *et al.*, 2002; Rodríguez *et al.*, 2006). On the other
26
27 298 hand, both strains have shown different degree of virulence and host specificity (Novoa *et al.*
28
29 299 1998; Alam *et al.* 2002). For the 56 possible similar proteins between both strains,
30
31 300 confirmation of identity was obtained in 80% of the selected spots, and only 3 proteins that
32
33 301 matched in position had a different identity. The 20 most abundant proteins were the same
34
35 302 between both *V. tapetis* strains, confirming that these proteins either not change in their amino
36
37 303 acid sequence between these strains, or present only minor changes that do not alter the pI of
38
39 304 the protein. This finding suggests that these abundant proteins are important for the proper
40
41 305 performance and survival of any strain of *V. tapetis*.

42
43
44 306
45
46
47
48 307 It is interesting to point out that the most abundant protein in both strains rendered no positive
49
50 308 match by MS or MS/MS, although by *de novo* sequencing it could be identified as Porin-like
51
52 309 protein H (OmpH). This finding confirms that both strains present the same most abundant
53
54 310 protein, even though showing slight changes in the amino acid sequence that causes an
55
56
57
58
59
60

1
2
3
4 311 alteration in the mobility of the protein at pI but not at molecular weight level. This protein is
5
6 312 one of the major outer membrane proteins and is present in very high copy numbers in the
7
8 313 bacterial surface (Lee *et al.* 2006).
9

10 314

11
12 315 On the basis of the identification results, it seems clear that, under the culture conditions
13
14 316 employed in this study, glucose degradation is one of the major pathways for energy
15
16 317 production, this hypothesis is supported by the fact that four enzymes, involved in four of the
17
18 318 ten steps of glycolysis, are among the 20 most abundant identified proteins. In addition, the
19
20 319 presence of elongation factors within these most abundant proteins indicates that *V. tapetis*
21
22 320 synthesizes and processes a great amount of proteins under the laboratory conditions
23
24 321 employed.
25
26
27

28 322 The reference map obtained in this study not only demonstrates the similarity under
29
30 323 laboratory conditions of two strains of the species with remarkable differences at genetic and
31
32 324 pathophysiological levels, but also could constitute the first basic step for a further
33
34 325 determination of important proteins and pathways involved in the virulence of *V. tapetis*.
35
36

37 326

38 39 327 **Acknowledgements**

40
41 328 This work was supported in part by grants AGL2006-13208-C02-01 and AGL2010-18438
42
43 329 from the Ministerio de Ciencia y Tecnología (Spain). J.B.C. and S.B. acknowledge the Xunta
44
45 330 de Galicia and the Ministerio de Ciencia e Innovación (MICINN), (Spain) for research
46
47 331 fellowships.
48
49

50 332
51
52
53
54
55
56
57
58
59
60

333 **REFERENCES**

- 334 Allam, B. and Ford, S.E. (2006) Effects of the pathogenic *Vibrio tapetis* on defence factors of
335 susceptible and non-susceptible bivalve species. I. Haemocyte changes following in vitro
336 challenge. *Fish Shellfish Immunol* **20**: 374-83.
- 337 Allam, B., Ashton-Alcox, K. and Ford, S.E. (2001) Hemocyte activities associated with
338 resistance to brown ring disease in *Ruditapes* spp. clams. *Dev Comp Immunol* **25**: 365-75.
- 339 Allam, B., Paillard, C. and Ford, S.E. (2002) Pathogenicity of *Vibrio tapetis*, the etiological
340 agent of Brown Ring Disease (BRD) in clams. *Dis. Aquat. Org.* **48**: 221-31.
- 341 Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W. and D.J.
342 Lipman. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database
343 search programs. *Nucleic Acids Res.* **25**:3389-3402.
- 344 Borrego, J.J., Castro, D., Luque, A., Paillard, C., Maes, P., García, M.T. and Ventosa, A.
345 (1996) *Vibrio tapetis* sp. nov., the causative agent of the brown ring disease affecting
346 cultured clams. *Int J Syst Bacteriol.* **46**:480-4.
- 347 Bumann, D. (2010) Pathogen proteomes during infection: A basis for infection research and
348 novel control strategies. *J. Proteomics.* **73**: 2267-76.
- 349 Cash, P. (2009) Proteomics in the study of the molecular taxonomy and epidemiology of
350 bacterial pathogens. *Electrophoresis* **30**:133-141.
- 351 Cash, P. and E. Argo. (2009) Analysis of bacterial proteins by 2DE. *Methods Mol. Biol.* **519**:
352 131-44.
- 353 Castro, D., Santamaria, J.A., Luque, A., Martínez-Manzanares, E. and Borrego J.J. (1996)
354 Antigenic characterization of the etiological agent of the brown ring disease affecting
355 Manila clams. *System. Appl. Microbiol.* **19**: 231-9.

- 1
2
3
4 356 Castro, D., Santamaría, J.A., Luque, A., Martínez-Manzanares, E. and J.J. Borrego (1997)
5
6 357 Determination of the etiological agent of brown ring disease in southwestern Spain. *Dis.*
7
8 358 *Aquat. Org.* **29**:181-8.
- 9
10 359 Choquet, G. (2004) Pathogénie de *Vibrio tapetis*, bacteria responsable de la maladie de
11
12 360 l'anneau brun chez la palourde: Approche cellulaire et moléculaire. Ph.D thesis. University
13
14 361 of Brest, France.
- 15
16 362 Choquet, G., Soudant, P., Lambert, C., Nicolas, J.L. and Paillard, C. (2003) Reduction of
17
18 363 adhesion properties on *Ruditapes philippinarum* hemocytes exposed to *Vibrio tapetis*. *Dis.*
19
20 364 *Aquat. Org.* **57**: 109-116.
- 21
22 365 Chung, W.J., Shu, H.Y., Lu C.Y., Wu, C.Y., Tseng, Y. H., Tsai, S.F. and Lin. C.H. (2007)
23
24 366 Qualitative and comparative proteomic analysis of *Xanthomonas campestris* pv.
25
26 367 *campestris* 17. *Proteomics.* **7**:2047-58.
- 27
28 368 Coelho, A., de Oliveira Santos, E., Faria, M.L., de Carvalho, D.P., Soares, M.R., von Kruger,
29
30 369 V.M. and Bisch, P.M. (2004) A proteome reference map for *Vibrio cholerae* El Tor.
31
32 370 *Proteomics.* **4**: 1491-1504.
- 33
34 371 De La Fuente, M., Borrajo, A., Bermúdez, J., Lores, M., Alonso, J., López, M., Santalla, M.,
35
36 372 De Ron, A.M., Zapata, C. and Alvarez G. (2011) 2-DE-based proteomic analysis of
37
38 373 common bean (*Phaseolus vulgaris* L.) seeds. *J. Proteomics.* **74**: 262-267.
- 39
40 374 Dunn, S.D., Revington, M., Cipriano, D.J. and Shilton, B.H. (2000) The b subunit of
41
42 375 *Escherichia coli* ATP synthase. *J. Bioenerg. Biomembr.* **32**:347-355.
- 43
44 376 Gietl, C. (1992) Malate dehydrogenase isoenzymes: cellular locations and role in the flow of
45
46 377 metabolites between the cytoplasm and cell organelles. *Biochim. Biophys. Acta.* **1100**:217-
47
48 378 234.
- 49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 379 Görg, A., Postel, W. and Günther, S. (1988) The current state of two-dimensional
5
6 380 electrophoresis with immobilized pH gradients. *Electrophoresis*. **9**:531-546.
7
8
9 381 Gragerov, A., Nudler, E., Komissarova, N., Gaitanaris, G.A., Gottesman, M.E. and Nikiforov,
10
11 382 V. (1992) Cooperation of GroEL/GroES and DnaK/DnaJ heat shock proteins in preventing
12
13 383 protein misfolding in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA*. **89**:10341-10344.
14
15 384 Hanson, R.W. and Patel, Y. M. (1994) Phosphoenolpyruvate carboxykinase (GTP): the gene
16
17 385 and the enzyme. *Adv. Enzymol. Relat. Areas Mol. Biol.* **69**: 203-281.
18
19 386 Hayano, T., Takahashi, N., Kato, S., Maki, N., Suzuki, M. (1991) Two distinct forms of
20
21 387 peptidylprolyl-cis-trans-isomerase are expressed separately in periplasmic and cytoplasmic
22
23 388 compartments of *Escherichia coli* cells. *Biochemistry*. **30**:3041-3048.
24
25
26 389 Jones, T.H., Johns, M.W. and Gill, C.O. (2008) Changes in the proteome of *Escherichia coli*
27
28 390 during growth at 15°C after incubation at 2, 6 or 8°C for 4 days. *Int. J. Food Microbiol.*
29
30 391 **124**: 299-302.
31
32
33 392 Kan, B., Habibi, H., Schmid, M., Liang, W., Wang, R., Wang, D. and Jungblut, P.R. (2004)
34
35 393 Proteome comparison of *Vibrio cholerae* cultured in aerobic and anaerobic conditions.
36
37 394 *Proteomics*. **4**: 3061-3067.
38
39
40 395 Kiraga, J., Mackiewicz, P., Mackiewicz, D., Kowalczyk, M., Biecek, P., Polak, N.,
41
42 396 Smolarczyk, K., Dudek, M.R. and Cebrat, S. (2007) The relationships between the
43
44 397 isoelectric point and: length of proteins, taxonomy and ecology of organisms. *BMC*
45
46 398 *Genomics* **8**:163.
47
48
49 399 Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of
50
51 400 bacteriophage T4. *Nature* **227**:680-685.
52
53
54
55
56
57
58
59
60

- 1
2
3
4 401 Lakhal, F., Bury-Moné, S., Nomane, Y., Le Goïc, N.L., Paillard, C. and Jacq, A. (2008) DjlA,
5
6 402 a membrane-anchored DnaJ-like protein, is required for cytotoxicity of clam pathogen *Vibrio*
7
8 403 *tapetis* to hemocytes. *Appl. Environm. Microbiol.* **74**: 5750-58.
- 9
10 404 Lane, E. and Birkbeck T.H. (2000) Species specificity of some bacterial pathogens of bivalve
11
12 405 molluscs is correlated with their interaction with bivalve haemocytes. *J. Fish Dis.* **23**: 275-
13
14 406 79.
- 15
16
17 407 Le Chevalier, P., Boula, C. and Paillard, C. (2003) Characterization by restriction fragment
18
19 408 length polymorphism and plasmid profiling of *Vibrio tapetis* strains. *J. Basic Microbiol.* **5**:
20
21 409 414–22.
- 22
23
24 410 Lee A,Y., Park, S.G., Jang, M., Cho, S., Myung, P.K., Kim, Y.R., Rhee, J.H., Lee, D.H and
25
26 411 Park, B.C. (2006) Proteomic analysis of pathogenic bacterium *Vibrio vulnificus*.
27
28 412 *Proteomics.* **6**:1283-1289.
- 29
30
31 413 López, J.R., Balboa, S., Núñez, S., de la Roca, E., de la Herran, R., Navas, J.I., Toranzo, A.E,
32
33 414 and Romalde, J.L. (2011) Characterization of *Vibrio tapetis* strains isolated from diseased
34
35 415 cultured Wedge sole (*Dicologlossa cuneata* Moreau). *Res. Vet. Sci.* **90**: 189-195.
- 36
37 416 López-Cortés, L., Luque, A., Martínez-Manzanares, E., Castro, D. and Borrego, J.J. (1999)
38
39 417 Adhesion of *Vibrio tapetis* to clam cells. *J. Shellfish Res.* **18**: 91-97.
- 40
41
42 418 Neuhoff, V., Arold, N., Taube, D. and Ehrhardt, W. (1988) Improved staining of proteins in
43
44 419 polyacrylamide gels including isoelectric focusing gels with clear background at nanogram
45
46 420 sensitivity using Coomassie Brilliant Blue G-250 and R-250. *Electrophoresis.* **9**: 255-262.
- 47
48 421 Nishimura, J.S., (1986) Succinyl-CoA synthetase structure-function relationships and other
49
50 422 considerations. *Adv. Enzymol. Relat. Areas Mol. Biol.* **58**:141-172.
- 51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 423 Novoa, B., Luque, A., Castro, D., Borrego, J.J. and Figueras, A. (1998). Characterization and
5
6 424 infectivity of four bacterial strains isolated from brown ring disease-affected clams. *J*
7
8 425 *Invertebr. Pathol.* **71**: 34-41.
- 9
10 426 Osman, K.M., Ali, M.M., Radwan, M. I., Kim, H.K. and Han, J. (2009) Comparative
11
12 427 proteomic analysis on *Salmonella gallinarum* and *Salmonella enteridis* exploring proteins
13
14 428 that may incorporate host adaptation in poultry. *J. Proteomics.* **72**: 815–821.
- 15
16
17 429 Paillard, C. (2004) A short-review of brown ring disease, a vibriosis affecting clams,
18
19 430 *Ruditapes philippinarum* and *Ruditapes decussatus*. *Aquat. Living. Resour.* **17**: 467-75.
- 20
21 431 Paillard, C. (2011) Vibriosis comparison between marine organisms models: is there
22
23 432 specificity in host-pathogen interactions? In J.L. Romalde (Ed) *Vibrio2011.The Fourth*
24
25 433 *Conference on the Biology of Vibrios.* pp. 93. Alvarellos SL, Lalín, Spain.
- 26
27
28 434 Paillard, C. and Maes, P. (1995) The brown ring disease in the Manila clam, *Ruditapes*
29
30 435 *philippinarum*. I. Ultrastructural alterations of the periostracal lamina. *J. Invertebr. Pathol.*
31
32 436 **65**: 91-100.
- 33
34
35 437 Paillard C, Maes, P. and Oubella. R. (1994) Brown ring disease in clams. *Annu. Rev. Fish*
36
37 438 *Dis.* **4**: 219-40.
- 38
39 439 Phillips, C.I. and Bogyo, M. (2005) Proteomics meets microbiology: technical advances in the
40
41 440 global mapping of protein expression and function. *Cell Microbiol.* **7**: 1061-76.
- 42
43
44 441 Poobalane, S., Thompson ,K.D., Diab, A., Ardó, L., Jeney, G. and Adams, A. (2008) Protein
45
46 442 expression by *Aeromonas hydrophila* during growth *in vitro* and *in vivo*. *Microb.*
47
48 443 *Pathogen.* **45**: 60-9.
- 49
50 444 Rodríguez, J.M., López-Romalde, S., Beaz, R., Alonso, M.C., Castro, D. and Romalde, J.L.
51
52 445 (2006) Molecular fingerprinting of *Vibrio tapetis* strains using three PCR-based methods:
53
54 446 ERIC-PCR, REP-PCR, and RAPD. *Dis Aquat Org* **69**:175-83.

- 1
2
3
4 447 Romalde, J.L., Castro, D., Magariños, B., Lopez-Cortes, L. and Borrego, J.J. (2002)
5
6 448 Comparison of ribotyping, randomly amplified polymorphic DNA, and pulsed-field gel
7
8 449 electrophoresis for molecular typing of *Vibrio tapetis*. *Syst. Appl. Microbiol.* **25**: 544-50.
9
10 450 Roncada, P., Deriu, F., Gaviraghi, A., Martino, P.A. and Bonizzi, L. (2009) Proteomic study
11
12 451 of antibiotic resistance in *Escherichia coli* strains. *Vet. Res. Commun.* **33**:157–60.
13
14 452 Sharma, P., Kumar, B. N., Singhal, V.M., Katoch, K., Venkatesan, D.S., Chauhan, D. and
15
16 453 Bisht D. (2010) Streptomycin induced protein expression analysis in *Mycobacterium*
17
18 454 *tuberculosis* by two-dimensional gel electrophoresis & mass spectrometry. *Indian J. Med.*
19
20 455 *Res.* **132**: 400-408.
21
22 456 Trinkler, N., Guichard, N., Labonne, M., Plasseraud, L., Paillard, C. and Marin. F. (2011)
23
24 457 Variability of shell repair in the Manila clam *Ruditapes philippinarum* affected by the
25
26 458 brown ring disease: a microstructural and biochemical study. *J. Invert. Pathol.* **106**:407-17.
27
28 459 Trinkler, N., Siquin, G., Querne, J. and Paillard, C. (2010) Resistance of brown ring disease
29
30 460 in the Manila clam, *Ruditapes philippinarum*: a study of selected stocks showing a
31
32 461 recovery process by shell repair. *J Invertebr. Pathol.* **104**: 8-16.
33
34 462 VanBogelen, R. A., Kelley, P. M .and Neidhardt, F. C. (1987) Differential induction of heat
35
36 463 shock, SOS, and oxidative stress regulons and accumulation of nucleotides in *Escherichia*
37
38 464 *coli*. *J. Bacteriol.* **169**:26-32.
39
40 465 von Krüger, W.M., Lery, L.M., Soares, M.R., de Neves-Manta, F.S., Batista e Silva, C.M.,
41
42 466 Neves-Ferreira, A.G., Perales, J. and Bisch, P.M. (2006) The phosphate-starvation
43
44 467 response in *Vibrio cholerae* O1 and *phoB* mutant under proteomic analysis: disclosing
45
46 468 functions involved in adaptation, survival and virulence. *Proteomics.* **6**:1495-511.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 471
5
6 472
7
8 473
9
10
11 474
12
13 475
14
15 476
16
17 477
18
19 478
20
21
22 479
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

480 **Table 1.** Proteins in the proteome of *Vibrio tapetis* CECT 4600^T identified by mass
 481 spectrometry.

482	COG Category	Protein	Spot	pI¹	Strain²
483	METABOLISM				
484	Carbohydrate transport and metabolism				
485		fructose-1,6-bisphosphatase	52, 149	5.6, 5.6	A/B
486		glucose-specific PTS system enzyme IIA component	49, 131	4.5, 4.5	A/B
487		aconitate hydratase	1, 82	4.9, 4.9	A/B
488		isocitrate dehydrogenase, NADP-dependent	4, 84	4.9, 4.9	A/B
489		succinyl-CoA synthetase	24, 39, 98,	4.9, 5.2, 4.8,	A/B
490			99, 109	4.9, 5.2	
491		succinate dehydrogenase iron-sulfur protein	43, 46	5.1, 5.3	A/B
492			116, 123	5.1, 5.3	
493		succinate dehydrogenase flavoprotein subunit	50, 152	5.5, 5.5	A/B
494		dihydrolipoamide dehydrogenase	51, 153	5.8, 5.7	A/B
495		enolase	18, 95	5.0, 5.1	A/B
496		fructose-bisphosphate aldolase	31, 101	4.9, 4.8	A/B
497		transaldolase B	35, 106	4.8, 4.8	A/B
498		glucose-1-phosphate adenylyltransferase	23	5.4	A
499		fumarate hydratase	11	5.2	A
500		citrate synthase	59	5.6	A
501		phosphoglycerate kinase	25*	4.9	A
502		glyceraldehyde-3-phosphate dehydrogenase	32	5.2	A
503		triosephosphate isomerase	42	4.9	A
504	Energy production and conversion				
505		malate dehydrogenase	34, 105	4.9, 4.9	A/B
506		Phosphoenolpyruvate carboxykinase	7, 86, 87	5.0, 4.9, 5.0	A/B
507		FOF1 ATP synthase subunit beta	9, 90	4.7, 4.7	A/B
508		ATPase	12, 94	5.2, 5.2	A/B
509		glycerol kinase	10	5.1	A
510		acetate kinase	21	5.2	A
511		deoxyribose-phosphate aldolase	113	4.9	B
512		isocitrate dehydrogenase, NADP-dependent	84	4.9	B
513					

514

515	COG Category	Protein	Spot	pI ¹	Strain ²
-----	--------------	---------	------	-----------------	---------------------

516 **Amino acid transport and metabolism**

517 serine hydroxymethyltransferase 22, 165, 163 5.3, 6.2, 6.1 A/B

518 alanine dehydrogenase 26, 28, 151 5.2, 5.3, 5.3 A/B

519 acetylornithine aminotransferase 29, 15 5.3, 4.9 A/B

520 amino acid ABC transporter protein 53, 146, 5.8, 6.0, A/B

521 148, 159 5.7, 6.7

522 aminopeptidase B 60, 154 5.8, 5.8 A/B

523 diaminopimelate decarboxylase 16 4.9 A

524 cysteine synthase A 33 5.3 A

525 ketol-acid reductoisomerase 8 4.6 A

526 aspartate aminotransferase 30 5.4 A

527 lysine/arginine/ornithine transporter subunit 147 5.8 B

528 **Nucleotide transport and metabolism**

529 uridine phosphorylase 44, 117 5.2, 5.2 A/B

530 purine nucleoside phosphorylase 76, 122 5.1, 5.0 A/B

531 adenylosuccinate synthetase 13 5.3 A

532 nucleoside diphosphate kinase 70 5.3 A

533 UDP-glucose/GDP mannose dehydrogenase 93 5.1 B

534 adenylate kinase 114 4.9 B

535 **Lipid metabolism**

536 3-hydroxydecanoyl-ACP dehydratase 65, 139 5.4, 5.4 A/B

537 **Coenzyme metabolism**

538 6,7-dimethyl-8-ribityllumazine synthase 68 5.5 A

539 uroporphyrinogen decarboxylase 164 5.8 B

540

541 **CELLULAR PROCESSES**542 **Posttranslational modification, protein turnover, chaperones**

543 molecular chaperone DnaK 3, 85 4.9, 4.9 A/B

544 heat shock protein 60 family chaperone GroEL 6, 89 4.8, 4.8 A/B

545 peptidyl-prolyl cis-trans isomerase 40, 120, 128 4.5, 4.5, 4.8 A/B

546 alkyl hydroperoxyde C22 protein 41, 118, 119 4.6, 4.8, 4.6 A/B

547 ATP-dependent Clp protease, subunit ClpP 74, 134 5.1, 5.1 A/B

548 DNA-binding H-NS 78, 136 5.1, 5.1 A/B

549 stringent starvation protein A 77 5.1 A

550	stress-induced protein UspE	81	5.6	A	
551					
552	COG Category	Protein	Spot	pI¹	Strain²
553	Signal transduction mechanisms				
554	dnaK supressor protein	73, 135	5.1, 5.1	A/B	
555	S-ribosylhomocysteinase	74	5.1	A	
556	stress-induced protein UspE	81	5.6	A	
557	cAMP regulatory protein	160	6.8	A	
558	Inorganic ion transport and metabolism				
559	superoxide dismutase [Fe]	48, 133	5.1, 5.1	A/B	
560	TRAP transporter solute receptor TAXI family precursor	157, 158	6.7, 6.8	A	
561	exopolyphosphatase like protein	110	5.0	B	
562	Cell envelope biogenesis, outer membrane				
563	outer membrane protein TolC	15, 97	4.9, 4.8	A/B	
564	Cell division and chromosome partitions				
565	chromosome segregation ATPase	124	5.4	B	
566					
567	INFORMATION STORAGE AND PROCESSING				
568	Translation, ribosomal structure and biogenesis				
569	elongation factor G	2, 83	4.9, 4.9	A/B	
570	elongation factor Tu	17, 125	4.9, 4.8	A/B	
571	50S ribosomal protein L10	56, 62,	6.1, 5.2,	A/B	
572		137, 144	5.2, 6.0		
573	30S ribosomal protein S1p	5, 57,	4.8, 6.0,	A/B	
574		88, 143	4.8, 6.0		
575	translation elongation factor Ts	38, 108	5.1, 5.1	A	
576	ribosome recycling factor	58	5.5	A	
577	elongation factor P	121	4.5	B	
578	tryptophanyl-tRNA synthetase	155	6.4	B	
579	tyrosyl-tRNA synthetase	162	6.0	B	
580	Transcription				
581	RNA polymerase alpha-subunit	25*, 100	4.9, 4.8	A/B	
582	transcription elongation factor GreA	79	4.6	A	
583	DNA replication, recombination and repair				
584	Holiday junction resolvase-like protein	27	5.2	A	

585

586

587

COG Category	Protein	Spot	pI ¹	Strain ²
POORLY CHARACTERIZED				
General function prediction only				
	hemolysin co-regulated protein Hcp-3	61	4.8	A
	isopenicillin N synthase	11	5.0	B
Function unknown				
	hypothetical protein pVT1_26	66, 140	5.6, 5.6	A/B
	hypothetical protein VSWAT3	45, 91	5.3, 4.7	A/B
	hypothetical protein VSAK1_11258	72	5.1	A
	hypothetical protein 1103602000595_AND4_07664	55	6.0	A
	putative type VI secretion protein VasQ-1	126	4.8	B
	hypothetical protein MED222	92, 129	4.8, 4.6	B
	UPF0065 protein in clcb-clcd intergenic region	156	6.4	B

600

601 Two proteins were identified in this spot.

602 ¹Estimated pI of the protein based on their position in the gel.603 ²A/B: protein was identified in both strains; A: protein was identified in CECT4600^T strain; B: protein was
604 identified in GR0202RD strain.

605

606

1
2
3
4 607 **Table 2.** Proteins in the proteomes of *Vibrio tapetis* CECT 4600^T and GR0202RD identified
5
6 608 by *de novo* secuencing.
7
8
9 609

SPOT	PROTEIN
14	Conserved hypothetical protein (<i>Vibrio parahaemolyticus</i>)
15	Type I secretion outer membrane protein TolC (<i>Vibrio harveyi</i>)
20	Beta-lactamase / Glucose-6-phosphate dehydrogenase (<i>Shewanella halifaxiensis</i> / <i>Streptococcus pneumoniae</i>)
26	No identification
37	Porin-like protein H (OmpH) (<i>Vibrio shilonii</i>)
54	No identification
55	hypothetical protein V12G01_08700 (<i>Vibrio alginolyticus</i>)
64	Glycine cleavage system regulatory protein (<i>Vibrio parahaemolyticus</i>)
67	Hypothetical protein VCG_002854 (<i>Vibrio cholerae</i>)
69	Transcript regulator TetR family (<i>Vibrio parahaemolyticus</i>)
71	No identification
80	No identification
96	Elongation factor Tu (<i>Vibrio splendidus</i>)
104	Porin-like protein H (OmpH) (<i>Vibrio shilonii</i>)

610

611

612 **Table 3.** Twenty most abundant proteins in the *V. tapetis* CECT4600^T 2-DE reference map.

Protein ^a	Spot N ^o ^b	COG category ^c
unknow protein	37	Function unknow
elongation factor Tu	17	Translation, ribosomal structure and biogenesis
alkyl hydroperoxide reductase	41	Posttranslational modification, protein turnover, chaperones
phosphoenolpyruvate carboxykinase	7	Energy production and conversion
alanine dehydrogenase	28	Amino acid transport and metabolism
glyceraldehyde-3-phosphate dehydrogenase	32	Carbohydrate transport and metabolism
enolase	18	Carbohydrate transport and metabolism
molecular chaperone DnaK	3	Posttranslational modification, protein turnover, chaperones
fructose-bisphosphate aldolase	31	Carbohydrate transport and metabolism
F0F1 ATP synthase subunit beta	9	Energy production and conversion
heat shock protein 60 family chaperone GroEL	6	Posttranslational modification, protein turnover, chaperones
malate dehydrogenase	34	Energy production and conversion
phosphoglycerate kinase / RNA polymerase alpha-subunit	25	Carbohydrate transport and metabolism/transcription
ATPase	12	Energy production and conversion
succinyl-CoA synthetase subunit beta	24	Energy production and conversion
succinyl-CoA synthetase subunit alpha	39	Energy production and conversion
Holliday junction resolvase-like protein [<i>Vibrio</i> sp.]	27	Low score; <i>de novo</i> sequencing.
elongation factor G	2	Translation, ribosomal structure and biogenesis
translation elongation factor Ts	38	Translation, ribosomal structure and biogenesis
peptidyl-prolyl cis-trans isomerase B	61	Post-translational modification, protein turnover, chaperones

613 ^a Proteins are listed from most abundant to less abundant according to their expression; ^b Spot number from Fig.
614 2; ^c COG categories according to <http://www.ncbi.nlm.nih.gov/COG/old/palox.cgi?fun=all>

615

616 **Table 4.** Relation between proteins with matched position and their identification by MS.

617	618			619		
620	Spot matched in position		Same identification	Spot matched in position		Same identification
621	CECT 4600 ^T	GR0202RD	by MS	CECT 4600 ^T	GR0202RD	by MS
621	1	82	Yes	47	124	Yes
622	2	83	Yes	48	133	Yes
623	3	85	Yes	49	131	Yes
624	4	84	Yes	50	152	Yes
625	5	88	Yes	51	153	Yes
626	6	89	Yes	52	149	Yes
627	7	87	Yes	53	148	Yes
628	8	91	Yes	54	146	No
629	9	90	Yes	55	146	No
630	12	94	Yes	56	144	Yes
631	15	97	Yes	57	143	Yes
632	17	96	Yes	60	154	Yes
633	18	95	Yes	61	128	Yes
634	24	99	Yes	62	137	Yes
635	25	100	Yes	63	126	No
636	28	151	Yes	64	127	No
637	29	150	Yes	65	139	Yes
638	31	101	Yes	66	140	Yes
639	34	105	Yes	67	141	No
640	35	106	Yes	68	148	No
641	38	108	Yes	73	135	Yes
642	39	109	Yes	74	134	Yes
643	40	120	Yes	76	122	Yes
644	41	119	Yes	78	136	Yes
645	43	116	Yes	79	130	No
646	44	117	Yes	80	129	No
647	46	123	Yes			
648						
649						

1
2
3
4 650
5 651
6
7 652
8
9
10 653
11
12 654
13
14 655
15
16 656
17
18 657
19
20
21 658
22
23 659
24
25 660
26
27 661
28
29 662
30
31
32 663
33
34 664
35
36 665
37
38 666
39
40 667
41
42 668
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure legends


Figure 1. Comparison of 2-DE patterns of the cytoplasmic proteins in two cultures of *Vibrio tapetis* CECT 4600^T. IEF was carried out using IPG strips (24cm, pH 4-7). A 12,5% SDS gel stained with Commassie G-250 was used in the second dimension. Three replicate gels of each culture were made in order to assess inter- and intra-variability of the method.

Figure 2. Reference map of *Vibrio tapetis* CECT 4600^T. The identified proteins are labeled with spot numbers (from 1 to 81 and 165) and listed in Table 1.

Figure 3. Fuctional categories of the cytoplasmic proteins of *Vibrio tapetis* CECT 4600^T.

Figure 4. Reference map of *Vibrio tapetis* GR0202RD. The identified proteins are labeled with spot numbers (from 82 to 164) and listed in Table 1.


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



119x77mm (300 x 300 DPI)


Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



246x190mm (300 x 300 DPI)


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



150x93mm (300 x 300 DPI)

Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



254x186mm (300 x 300 DPI)

Review