



# Color coherence in a heavy quark antenna radiating gluons inside a QCD medium



Manoel R. Calvo<sup>a,b</sup>, Manoel R. Moldes<sup>a,c,\*</sup>, Carlos A. Salgado<sup>a</sup>

<sup>a</sup> Departamento de Física de Partículas and IGFAE, Universidade de Santiago de Compostela, 15706 Santiago de Compostela, Galicia, Spain

<sup>b</sup> Institut de Physique Théorique, Saclay, France

<sup>c</sup> Centre de Physique Théorique, École Polytechnique, CNRS, 91128 Palaiseau, France

## ARTICLE INFO

### Article history:

Received 9 May 2014

Received in revised form 19 September 2014

Accepted 4 October 2014

Available online 8 October 2014

Editor: J.-P. Blaizot

### Keywords:

Perturbative QCD

Jet quenching

Heavy quarks

## ABSTRACT

We compute the color coherence effects for soft gluon radiation off antennas containing heavy quarks in the presence of a QCD medium – the actual calculations is made for a triplet configuration and then generalize to both color singlet and octet ones. This work completes the studies of antenna radiation inside a medium which provide a useful picture of the relevance of interference effects in jet parton showers for the jet quenching phenomenon observed in high-energy nuclear collisions. The analysis is performed resumming the multiple scatterings of the partonic system with the medium. The main conclusion is that decorrelation due to color rotation is more effective in the case in which at least one of the emitters of the antenna is a heavy quark. This effect, present both for a heavy-quark-antiquark or a heavy-quark-gluon antenna is more relevant for the later or for the case in which the energies of the quark and antiquark are very different. The parameter controlling these effects involves the dead-cone angle. We find that interferences are cancelled, spoiling the color correlation of the pair, when  $\theta_{DC} \equiv M/E \gg 1/\sqrt{\omega L}$  where  $E$  and  $\omega$  are the energies of the heavy quark and the radiated gluon and  $L$  is the medium length. In the case of a heavy-quark-antiquark antenna  $t_{\text{form}}$ , defined as the difference in splitting times in amplitude and complex conjugate of the amplitude, appears instead of  $L$  if the original splitting is symmetric. The presence or absence of interferences modifies the energy loss pattern.

© 2014 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>). Funded by SCOAP<sup>3</sup>.

## 1. Introduction

Jets traversing QCD matter created in high-energy nuclear collisions have been experimentally studied for the last ten years, first at the RHIC at BNL [1] and then at the LHC at CERN [2–4]. Despite the success of a theory of jet quenching based on an enhancement of the gluon radiation induced by the medium [5–12] (for a recent review see e.g. [13]) mainly for the description of the data on inclusive particle suppression, a complete theory, suitable for a consistent and rigorous interpretation of the reconstructed jet data, is still being developed. On the theoretical side, progress in the last few years has been reached in different fronts, as improvements in the splitting probability [14,15], the use of effective theories as SCET [16–18], or the study of multi-parton radiation [17]. A systematic program to understand the in-medium intra-

jet color coherence effects, using the *antenna* setup, has also been started, leading to extremely valuable information about the role of interferences [24–30]. A clear new picture of the jet quenching phenomena is emerging from these studies [31,32].

Since the antenna spectrum plays a central role in understanding jets physics in vacuum [19–23], it seems quite natural to ask about the case in what the antenna is traversing a dense medium. The basic question which is addressed with the antenna setup is to which extent subsequent gluon emissions can be considered as independent, hence providing a clear probabilistic picture, and how and when this independency is broken. The well-known results of color coherence in the vacuum lead to the picture of angular-ordered emissions of gluons. The picture in the medium follows the same basic principles and can be simplified as follows [31]: medium-induced radiation can only resolve objects (emitters) which are separated more than a transverse distance  $\Lambda_{\text{med}}$  determined by the medium properties. Taking the example of a quark-antiquark antenna, when the transverse separation  $r_{\perp} \sim \theta_{q\bar{q}}L \ll \Lambda_{\text{med}}$  the medium cannot resolve the quark and the antiquark

\* Corresponding author at: Departamento de Física de Partículas and IGFAE, Universidade de Santiago de Compostela, 15706 Santiago de Compostela, Galicia, Spain.  
E-mail address: [manoel.rodriguez-moldes@usc.es](mailto:manoel.rodriguez-moldes@usc.es) (M.R. Moldes).

individually, so they remain color correlated and emit coherently. This emission can be decomposed into medium-induced (soft and collinear finite) radiation by the total charge, i.e. no radiation if the pair is in a singlet state or radiation as a gluon if the pair is in octet state, plus a vacuum like component which is soft and collinear divergent and angular ordered. In the opposite case  $r_{\perp} \sim \theta_{q\bar{q}} L \gg \Lambda_{\text{med}}$  the medium is very efficient in destroying the color correlation of the pair and the quark and antiquark emit incoherently. Two components can be also distinguished, two medium-induced (soft and collinear finite) contributions each with strength  $C_F$  as it corresponds to a singlet emitter, plus a vacuum-like radiation (soft and collinear divergent) but where angular ordering is removed, as expected from two independent color sources.

The mass of the heavy quarks is known to modify the role of color coherence, e.g. introducing a dead-cone angle where radiation is strongly suppressed or removing the strict angular ordering in the vacuum. In the case of the medium, similar modifications were observed some time ago for the case of a single emitter [33–36]: on the one hand, a suppression of the radiation was predicted in most of the phase space relevant for the phenomenological applications; on the other hand, the smaller typical formation time of the gluons produced off massive quarks lead to a reduction of the Landau–Pomeranchuk–Migdal suppression enhancing the radiation at small angles which to some extent fills the dead cone [34]. This additional radiation is not very relevant for the total energy loss of heavy quarks which turns out to be smaller than for light quarks – this is so because it happens only in a reduced region of phase space, of little phenomenological relevance [34]. The corresponding experimental search of this *dead-cone effect* lead to one of the still unsolved puzzles in RHIC data [37,38] in which a suppression of the non-photon electrons (expected to be dominated by heavy-quark decays) is compatible, taking at face value, with no mass-effect in the radiation. Recent LHC results seem to indicate that indeed heavy-quarks lose less energy than light partons [39–41] but the actual effect still needs to be quantified as several different mechanisms contribute to the observed suppression of heavy mesons – e.g. the different slopes of the perturbative spectra or the harder fragmentation functions in the case of heavy quarks. Interesting results also exist for the case of b-quark jets [42] which study a region of much larger transverse momentum, where the effects of the mass are expected to be small – some coherence effects as the ones studied here are, however, expected to be of relevance there.

Motivated by these theoretical and experimental findings we present here a calculation of the color coherence effects in a set up which includes a heavy-quark and a gluon antenna radiating a soft gluon in the presence of a medium. The calculation is done resumming the multiple scatterings of the partons involved with the surrounding medium. We also comment on the case of a heavy-quark–antiquark antenna, previously studied in Ref. [27] but only for the case of the first order in the opacity expansion. The main result in the paper is the reduction of the color coherence when the dead-cone angle  $\theta_{DC} \equiv M/E \gg 1/\sqrt{\omega L}$  where  $E$  and  $\omega$  are the energies of the heavy quark and the radiated gluon and  $L$  is the medium length.<sup>1</sup> This suppression of the color coherence enhances the phase space for radiation, hence the energy loss of the heavy quark. Whether this enhancement can lead to a sizable suppression of the heavy quarks in the experimental environment is left for a future study. The present publication completes the study of the in-medium antenna radiation spectrum [24–30] and provides

the way to encode heavy quark effects in the new picture of jet quenching being developed.

## 2. Amplitudes and formalism

The derivation of the spectrum is similar to the ones presented in previous works [24–30] but including the mass of the heavy quark explicitly. The amplitude for one gluon emission is calculated using the reduction formula

$$\mathcal{M}^a(k) = - \sum_{\lambda} \int_{x^+=+\infty} dx^- d^2\mathbf{x} e^{ik \cdot x} 2\partial_x^+ \mathbf{A}^a(x) \cdot \boldsymbol{\epsilon}_{\lambda}(\vec{k}) \quad (1)$$

with  $k^{\mu} = (\omega, \vec{k})$  being the 4-momentum of the emitted gluon and  $\mathbf{A}$  the transverse gauge field. The gauge field is obtained from the classical Yang–Mills (CYM) equations

$$[D_{\mu}, F^{\mu\nu}] = J^{\nu} \quad (2)$$

where  $D_{\mu} \equiv \partial_{\mu} - igA_{\mu}$  and  $F_{\mu\nu} \equiv \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu} - ig[A_{\mu}, A_{\nu}]$ , and with the current  $J^{\mu}$  being covariantly conserved, i.e.,

$$[D_{\mu}, J^{\mu}] = 0 \quad (3)$$

The current  $J^{\mu}$  has three components: one of them for each leg of the antenna and a third one representing the highly virtual particle coming from a hard process that splits into the antenna pair. For the case of a  $Qg$  antenna, this means that  $J^{\mu} = J_q^{\mu} + J_g^{\mu} + J_3^{\mu}$ , being the three components are the currents representing the quark, the gluon and the virtual quark, respectively.

The initial state of the antenna is given by the vacuum current  $J_{(0)}^{\mu} = J_{q(0)}^{\mu} + J_{g(0)}^{\mu} + J_{3(0)}^{\mu}$ , where

$$J_{i(0)}^{\mu,a}(x) = g \frac{p_i^{\mu}}{E_i} \delta^{(3)}\left(\vec{x} - \frac{\vec{p}_i}{E_i} t\right) \theta(t) Q_i^a \quad (4)$$

represents a particle with momentum  $p_i^{\mu} = (E_i, \vec{p}_i)$  and charge color vector  $Q_i^a$  and  $i = q, g, 3$ .

The  $J_3^{\mu}$  current is needed for charge and momentum conservation ( $Q_3 = Q_q + Q_g$  and  $\vec{p}_3 = -\vec{p}_q - \vec{p}_g$ , respectively), but in a colored antenna it does not contribute in the frame where  $p_3^{\mu} \simeq (0, p_3^-, \mathbf{0})$  due to the light-cone gauge we perform our calculation in ( $A^+ = 0$ ).<sup>2</sup> In the case of a singlet antenna,  $J_3^{\mu}$  does not contribute because  $Q_3 = 0$ .

The effect of the medium over the vacuum current  $J_{(0)}^{\mu}$  is to induce a color rotation:

$$J^{\mu}(x) = U_p(x^+, 0) J_{q(0)}^{\mu}(x) + U_{\bar{p}}(x^+, 0) J_{g(0)}^{\mu}(x) \quad (5)$$

described by a Wilson line:

$$U_p(x^+, 0; \mathbf{r}) \equiv \mathcal{P} \exp \left\{ \int_0^{x^+} d\xi T \cdot A_{\text{med}}^-(\xi, \mathbf{p}\xi/p^+) \right\} \quad (6)$$

where we have denoted the quark momentum as  $p$  and the gluon momentum as  $\bar{p}$ .

Leaving only terms linear on the medium induced field and focusing on the quark current (the calculation with the gluon current is the same and so it gives an analogous result) we get the following expression for the amplitude for emission off the quark:

<sup>1</sup> Notice that in this paper, light-cone coordinates are used, so, the definition of the dead-cone angle  $\theta_{DC}$  will be slightly different, see below.

<sup>2</sup> Any 4-vector  $a^{\mu} = (a^0, a^1, a^2, a^3)$  is expressed in light-cone coordinates as  $a^{\mu} = (a^+, a^-, \mathbf{a})$ , with  $a^{\pm} = (a^0 \pm a^3)/\sqrt{2}$  and  $\mathbf{a} = (a^1, a^2)$ .

$$\begin{aligned} \mathcal{M}_q^a(\vec{k}) &= \sum_{\lambda} \frac{g}{k^+} \int_{x^+=+\infty} d^2\mathbf{x} e^{ik^-x^+} e^{-i\mathbf{k}\cdot\mathbf{x}} \int_0^{+\infty} dy^+ e^{ik^+ \frac{p^-}{p^+} y^+} \\ &\times \epsilon_{\lambda}(k) \cdot (i\partial_y + k^+ \mathbf{n}) \\ &\times \mathcal{G}^{ab}(x^+, \mathbf{x}; y^+, \mathbf{y}|k^+) \Big|_{\mathbf{y}=\mathbf{n}y^+} U_p^{bc}(y^+, 0) Q_q^c \end{aligned} \quad (7)$$

where we have explicitly stated the color structure, defined the dimensionless vector  $\mathbf{n} = \mathbf{p}/p^+$  and  $\mathcal{G}$  is a Green's function that takes into account both the color rotation of the gluon and its Brownian motion in the transverse plane due to interactions with the medium field. These features of the Green's function  $\mathcal{G}$  can be easily seen thanks to its expression as a path integral in the transverse plane:

$$\begin{aligned} \mathcal{G}(x^+, \mathbf{x}; y^+, \mathbf{y}|k^+) &= \int_{\mathbf{r}(y^+)=\mathbf{y}}^{\mathbf{r}(x^+)=\mathbf{x}} \mathcal{D}\mathbf{r} \exp \left\{ \frac{ik^+}{2} \int_{y^+}^{x^+} d\xi \dot{\mathbf{r}}^2(\xi) \right\} U(x^+, y^+; \mathbf{r}) \end{aligned} \quad (8)$$

The mass effects can be now easily identified, as they enter the ‘-’ component of the 4-momentum heavy-quark momentum through the dispersion relation

$$2p^+ p^- - \mathbf{p}^2 = M^2. \quad (9)$$

Taking the eikonal limit, in which the quark follows a straight line in the direction  $\mathbf{n}$ , the presence of the mass appears only in the phase

$$\exp \left( ik^+ \frac{p^-}{p^+} y^+ \right) = \exp \left( i \frac{k^+}{2} \theta_{DC}^2 y^+ \right) \exp \left( i \frac{k^+}{2} \mathbf{n}^2 y^+ \right) \quad (10)$$

where  $\theta_{DC}$  is the so-called *dead-cone angle*, defined as<sup>3</sup>

$$\theta_{DC} \equiv \frac{M}{p^+} \quad (11)$$

We have now all the ingredients to explicitly write the two contributions to the amplitude; when the gluon is radiated off the heavy quark

$$\begin{aligned} \mathcal{M}_q^a(\vec{k}) &= \sum_{\lambda} \frac{g}{k^+} \int_{x^+=+\infty} d^2\mathbf{x} e^{ik^-x^+} e^{-i\mathbf{k}\cdot\mathbf{x}} \\ &\times \int_0^{+\infty} dy^+ \exp \left[ i \frac{k^+}{2} (\theta_{DC}^2 + \mathbf{n}^2) y^+ \right] \\ &\times \epsilon_{\lambda}(k) \cdot (i\partial_y + k^+ \mathbf{n}) \\ &\times \mathcal{G}^{ab}(x^+, \mathbf{x}; y^+, \mathbf{y}|k^+) \Big|_{\mathbf{y}=\mathbf{n}y^+} U_p^{bc}(y^+, 0) Q_q^c \end{aligned} \quad (12)$$

and when the gluon is radiated off the gluon

$$\begin{aligned} \mathcal{M}_g^a(\vec{k}) &= \sum_{\lambda} \frac{g}{k^+} \int_{x^+=+\infty} d^2\mathbf{x} e^{ik^-x^+} e^{-i\mathbf{k}\cdot\mathbf{x}} \int_0^{+\infty} dy^+ \exp \left[ i \frac{k^+}{2} \bar{\mathbf{n}}^2 y^+ \right] \\ &\times \epsilon_{\lambda}(k) \cdot (i\partial_y + k^+ \bar{\mathbf{n}}) \\ &\times \mathcal{G}^{ab}(x^+, \mathbf{x}; y^+, \mathbf{y}|k^+) \Big|_{\mathbf{y}=\bar{\mathbf{n}}y^+} U_p^{bc}(y^+, 0) Q_g^c \end{aligned} \quad (13)$$

We can explicitly see the difference due to the presence of mass: a complex phase in (12) that (13) lacks.

<sup>3</sup> Notice that this definition is slightly different that the one given in Section 1. Translating the light-cone coordinates into space ones,  $p^+ \simeq \sqrt{2}E$ , the dead-cone angle reads  $\theta_{DC} \simeq M/(\sqrt{2}E)$ .

### 3. Radiation spectrum of the heavy-quark–gluon antenna

With the amplitudes (12) and (13) we can compute the radiation spectrum of the antenna. A particularly convenient way of presenting this spectrum is by separating the independent radiation (in the vacuum easily identified by the two collinear divergencies) of the two emitters

$$dN = \frac{\alpha_s}{(2\pi)^2} [C_F \mathcal{R}_q + C_A \mathcal{R}_g - C_A \mathcal{J}] \frac{d^3k}{(k^+)^3} \quad (14)$$

Here, we have defined the independent radiation off the heavy quark  $\mathcal{R}_q$  by

$$C_F \mathcal{R}_q = (k^+)^2 |\mathcal{M}_q|^2 \quad (15)$$

the independent radiation off the gluon  $\mathcal{R}_g$

$$C_A \mathcal{R}_g = (k^+)^2 |\mathcal{M}_g|^2 \quad (16)$$

and the interference spectrum between both emitters  $\mathcal{J}$

$$-C_A \mathcal{J} = (k^+)^2 \text{Re} \langle \mathcal{M}_q \mathcal{M}_g^\dagger \rangle \quad (17)$$

The spectrum of independent radiation off the heavy quark  $\mathcal{R}_q$  is directly evaluated from (15) taking the limit  $\mathbf{n} \rightarrow 0$

$$\begin{aligned} \mathcal{R}_q &= 2 \text{Re} \int_0^\infty dy'^+ \int_0^{y'^+} dy^+ \exp \left[ i \frac{k^+}{2} \theta_{DC}^2 (y^+ - y'^+) \right] \\ &\times \int d^2\mathbf{z} \exp \left[ -i\mathbf{k} \cdot \mathbf{z} - \frac{1}{2} \int_{y'^+}^\infty d\xi n(\xi) \sigma(\mathbf{z}) \right] \\ &\times \partial_y \cdot \partial_z \mathcal{K}(y'^+, \mathbf{z}; y^+, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{0}} \end{aligned} \quad (18)$$

which, as expected, is exactly the same expression derived previously for the medium-induced gluon radiation off a single heavy quark in the BDMPs multiple scattering approximation [34]. This confirmation is also a test of our formalism.

In the same manner, the medium-induced gluon radiation off a gluon is obtained from (16)

$$\begin{aligned} \mathcal{R}_g &= 2 \text{Re} \int_0^\infty dy'^+ \int_0^{y'^+} dy^+ \\ &\times \int d^2\mathbf{z} \exp \left[ -i\mathbf{k} \cdot \mathbf{z} - \frac{1}{2} \int_{y'^+}^\infty d\xi n(\xi) \sigma(\mathbf{z}) \right] \\ &\times \partial_y \cdot \partial_z \mathcal{K}(y'^+, \mathbf{z}; y^+, \mathbf{y}) \Big|_{\mathbf{y}=\mathbf{0}} \end{aligned} \quad (19)$$

which again coincides with the known results.

The most interesting part of our analysis is, of course, the color coherent emission off the two emitters given by the interference term  $\mathcal{J}$

$$\begin{aligned} \mathcal{J} &= \text{Re} \left\{ \int_0^\infty dy'^+ \int_0^{y'^+} dy^+ (1 - \Delta_{\text{med}}(y^+, 0)) \right. \\ &\times \exp \left[ i \frac{k^+}{2} (\theta_{DC}^2 + \delta \mathbf{n}^2) y^+ \right] \\ &\times \int d^2\mathbf{z} \exp \left[ -i\vec{k} \cdot \mathbf{z} - \frac{1}{2} \int_{y'^+}^\infty d\xi n(\xi) \sigma(\mathbf{z}) \right] \end{aligned}$$

$$\times (\partial_y - ik^+ \delta \mathbf{n}) \cdot \partial_z \mathcal{K}(y'^+, \mathbf{z}; y^+, \mathbf{y}) \Big|_{\mathbf{y}=\delta \mathbf{n} y^+} \Big\} + \text{sym.} \quad (20)$$

with  $\bar{\kappa} = \mathbf{k} - k^+ \bar{\mathbf{n}}$  (and  $\kappa = \mathbf{k} - k^+ \mathbf{n}$ ),  $\delta \mathbf{n} = \mathbf{n} - \bar{\mathbf{n}}$  and  $\mathcal{K}$  being the path integral

$$\mathcal{K}(y'^+, \mathbf{z}; y^+, \mathbf{y} | k^+) = \int_{\mathbf{r}(y^+)=\mathbf{y}}^{\mathbf{r}(y'^+)=\mathbf{z}} \mathcal{D}\mathbf{r} \exp \left\{ \int_{y^+}^{y'^+} d\xi \left( i \frac{k^+}{2} \dot{\mathbf{r}}^2(\xi) - \frac{1}{2} n(\xi) \sigma(\mathbf{r}) \right) \right\} \quad (21)$$

that takes into account the emitted gluon multiple scattering with the medium and its Brownian motion in the transverse plane from  $\mathbf{r}(y^+) = \mathbf{y}$  to  $\mathbf{r}(y'^+) = \mathbf{z}$ . The symmetric part is obtained exchanging  $q \leftrightarrow g$ .

Eq. (20) is the main result of this paper. Compared with the light-quark case, the only difference is the presence of a new phase including the dead-cone angle  $\theta_{DC}$ . This is similar to the case found for the single heavy quark emitter [34] and a direct consequence of the eikonal approximation assumed in the calculation. The role of the phase is to suppress the radiation in some regions of phase space due to the mass effects.

#### 4. Discussion and conclusions

We will focus here in the two main terms which suppress the interferences, Eq. (20), namely the decoherence parameter  $\Delta_{\text{med}}$ :

$$\Delta_{\text{med}}(y^+, 0) \equiv 1 - \exp \left[ -\frac{1}{2} \int_0^{y^+} d\xi n(\xi) \sigma(\delta \mathbf{n} \xi) \right] \simeq 1 - \exp \left[ -\frac{1}{12} \hat{q} \delta \mathbf{n}^2 L^3 \right] \quad (22)$$

(where we have used the multiple soft scattering approximation for the last expression,  $\hat{q}$  being the transport coefficient also known as *jet quenching parameter*) and the *dead-cone* phase:

$$\Delta_{DC}(y^+, y'^+) = \exp \left[ i \frac{k^+}{2} (\theta_{DC}^2 y^+ - \bar{\theta}_{DC}^2 y'^+) \right] \quad (23)$$

Notice that for a (triplet)  $Qg$  antenna,  $\bar{\theta}_{DC} = 0$  and we recover the phase in Eq. (20). For an octet, or singlet,  $Q\bar{Q}$ , there are two dead-cone angles, each one referring to the energy of the corresponding emitter,  $\theta_{DC} = M/p^+$  and  $\bar{\theta}_{DC} = M/\bar{p}^+$ . Notice also that in the  $Q\bar{Q}$  case, when one of the emitters is much more energetic than the other, say  $\bar{p}^+ \gg p^+$ , i.e. for very asymmetric  $g \rightarrow Q\bar{Q}$  splittings, the suppression pattern of the interferences is similar to that of the  $Qg$  case, as  $\theta_{DC} \ll \bar{\theta}_{DC}$  in (23).

The effect of the decoherence parameter  $\Delta_{\text{med}}$  in the in-medium antenna radiation has been discussed at length in the previous calculations, in particular in Ref. [31]. Its role is to suppress the interference terms (20) when the transverse size of the antenna is larger than the typical medium color correlation length in the transverse plane. I.e. when the correlation length is larger than the size of the pair,  $\Delta_{\text{med}} \rightarrow 0$ , the medium cannot resolve the individual emitters, which act as a single object with the total charge of the pair ( $C_F$  for triplet,  $C_A$  for octet or 0 for singlet). In the opposite case,  $\Delta_{\text{med}} \rightarrow 1$ , the medium resolves the antenna and breaks the color coherence of the pair so that they behave as two independent particles. One way of estimating the effects of coherence is by defining a coherence time  $t_{\text{coh}} \sim [12/(\theta_{qq}^2 \hat{q})]^{1/3}$ , see Eq. (22), so that the pair stays coherent (acting as a single emitter) for medium path lengths of the antenna smaller than  $t_{\text{coh}}$ .

Of all three antennas involving heavy quarks (singlet  $Q\bar{Q}$ , triplet  $Qg$  and octet  $Q\bar{Q}$ ), the case of a singlet antenna can be safely neglected in all calculations of energy loss. The other two cases, triplet and octet, need to be considered and are relevant for different regions of the phase space. In particular, the octet  $Q\bar{Q}$  antenna is relevant for the case of high- $p_T$  heavy quarks produced by the decay of boosted virtual gluons.

So, we consider now the case of a color octet  $Q\bar{Q}$  antenna in the symmetric case ( $p^+ \simeq \bar{p}^+$ ) and assuming that the opening angle  $\delta \mathbf{n}^2 \sim \theta_{qq}^2$  is small (which is a comfortable assumption  $\theta_{qq}^2 \ll 1/(k^+L)$ ) and can be neglected in the phase. When color decoherence happens,  $\Delta_{\text{med}} \sim 1$ , the radiation is that of two independent emitters

$$(k^+)^3 \frac{dN}{d^3k} = \frac{\alpha_s}{(2\pi)^2} C_F (\mathcal{R}_q + \mathcal{R}_{\bar{q}}) \quad (24)$$

and the spectrum is just the superposition of two spectra for gluon radiation off a heavy-quark. This behavior was first identified for the massless case in [25].

An interesting case appears in the totally coherent case,  $\Delta_{\text{med}} \sim 0$ , in which the medium cannot resolve the pair, and one has  $\mathcal{R}_q \sim \mathcal{R}_{\bar{q}} \sim \mathcal{J}$ . Now, the medium-induced radiation is simply

$$(k^+)^3 \frac{dN}{d^3k} = \frac{\alpha_s}{(2\pi)^2} C_A \mathcal{R}_q \quad (25)$$

That would correspond to the radiation *off a gluon* (as the total charge gives the factor  $C_A$ ) *but with the dead-cone suppression factor*. As mentioned above, the production of heavy quarks at high transverse momentum receives a large contribution from the splitting of boosted gluons into a  $Q\bar{Q}$  pair. Before losing coherence this constitutes an octet  $Q\bar{Q}$  antenna. In this case, the corresponding energy loss, from Eq. (25), will be as the one for gluons but with a dead-cone suppression as long as the pair stays in a color coherent state. The corresponding time for this to occur,  $t_{\text{coh}} \sim [12/(\theta_{qq}^2 \hat{q})]^{1/3}$  can be sizable according to the estimates in [31]. Notice that in this case, the total radiation is still larger than the sum of the radiation of the  $Q\bar{Q}$  pair (24) as  $2C_F < C_A$ . (For the rest of the color configurations, color coherence reduces the amount of energy loss.)

Another consequence of our analysis is that for the case of asymmetric splitting, either a  $Qg$  antenna or a  $Q\bar{Q}$  antenna with very different energies of quark and antiquark, the suppression factor is much larger than for the symmetric case. Taking into account that  $k^+ \simeq \sqrt{2}\omega$ , it can be seen from Eq. (23) that for the symmetric configuration ( $\theta_{DC} \simeq \bar{\theta}_{DC}$ ) the suppression will happen for gluon energies<sup>4</sup>  $\omega > 1/(t_{\text{form}} \theta_{DC}^2)$ , while for the asymmetric case ( $\bar{\theta}_{DC} = 0$ ) it will happen much earlier,  $\omega > 1/(L\theta_{DC}^2)$ .

All these features provide a nice generalization of the findings of the antenna for the massless case with physics dominated by color coherence effects or the lost of them due to the interaction with the medium. The phenomenological consequences for the heavy quark case will be presented in a separate work.

#### Acknowledgements

This work is supported by European Research Council grant HotLHC ERC-2011-StG-279579, by Xunta de Galicia (Consellería de Educación); by the Spanish Consolider-Ingenio 2010 Programme CPAN and by FEDER. Manolo R. Calvo thanks Ministerio de Ciencia e Innovación of Spain for financial support under grant FPI-BES-2010-040744.

<sup>4</sup> The formation time is the difference in splitting times between the amplitude and the complex conjugate of the amplitude,  $t_{\text{form}} = y^+ - y'^+$ .

## References

- [1] K. Adcox, et al., PHENIX Collaboration, Nucl. Phys. A 757 (2005) 184; B.B. Back, et al., PHOBOS Collaboration, Nucl. Phys. A 757 (2005) 28; I. Arsene, et al., BRAHMS Collaboration, Nucl. Phys. A 757 (2005) 1; J. Adams, et al., STAR Collaboration, Nucl. Phys. A 757 (2005) 102.
- [2] K. Aamodt, et al., ALICE Collaboration, Phys. Lett. B 696 (2011) 30–39, arXiv:1012.1004 [nucl-ex].
- [3] G. Aad, et al., ATLAS Collaboration, Phys. Rev. Lett. 105 (2010) 252303, arXiv:1011.6182 [hep-ex].
- [4] S. Chatrchyan, et al., CMS Collaboration, Phys. Rev. C 84 (2011) 024906, arXiv:1102.1957 [nucl-ex].
- [5] R. Baier, Y.L. Dokshitzer, A.H. Mueller, S. Peigne, D. Schiff, Nucl. Phys. B 483 (1997) 291.
- [6] R. Baier, Y.L. Dokshitzer, A.H. Mueller, S. Peigne, D. Schiff, Nucl. Phys. B 484 (1997) 265.
- [7] B.G. Zakharov, JETP Lett. 63 (1996) 952.
- [8] B.G. Zakharov, JETP Lett. 65 (1997) 615.
- [9] U.A. Wiedemann, Nucl. Phys. B 588 (2000) 303.
- [10] M. Gyulassy, P. Levai, I. Vitev, Nucl. Phys. B 594 (2001) 371.
- [11] X.-N. Wang, X.-f. Guo, Nucl. Phys. A 696 (2001) 788, arXiv:hep-ph/0102230.
- [12] P.B. Arnold, G.D. Moore, L.G. Yaffe, J. High Energy Phys. 0206 (2002) 030, arXiv:hep-ph/0204343.
- [13] Y. Mehtar-Tani, J.G. Milhano, K. Tywoniuk, Int. J. Mod. Phys. A 28 (2013) 1340013, arXiv:1302.2579 [hep-ph].
- [14] L. Apolinario, N. Armesto, C.A. Salgado, Phys. Lett. B 718 (2012) 160, arXiv:1204.2929 [hep-ph].
- [15] J.-P. Blaizot, F. Dominguez, E. Iancu, Y. Mehtar-Tani, J. High Energy Phys. 1301 (2013) 143, arXiv:1209.4585 [hep-ph].
- [16] F. D'Eramo, M. Lekaveckas, H. Liu, K. Rajagopal, J. High Energy Phys. 1305 (2013) 031, arXiv:1211.1922 [hep-ph].
- [17] M. Fickinger, G. Ovanessian, I. Vitev, J. High Energy Phys. 1307 (2013) 059, arXiv:1304.3497 [hep-ph].
- [18] F. D'Eramo, H. Liu, K. Rajagopal, Phys. Rev. D 84 (2011) 065015, arXiv:1006.1367 [hep-ph].
- [19] A.H. Mueller, Phys. Lett. B 104 (1981) 161.
- [20] B.I. Ermolaev, V.S. Fadin, JETP Lett. 33 (1981) 269, Pis'ma Zh. Eksp. Teor. Fiz. 33 (1981) 285.
- [21] A. Bassetto, M. Ciafaloni, G. Marchesini, Phys. Rep. 100 (1983) 201; A. Bassetto, M. Ciafaloni, G. Marchesini, A.H. Mueller, Nucl. Phys. B 207 (1982) 189.
- [22] K. Konishi, A. Ukawa, G. Veneziano, Nucl. Phys. B 157 (1979) 45–107.
- [23] Y.L. Dokshitzer, V.A. Khoze, A.H. Mueller, S.I. Troyan, Basics of Perturbative QCD, Ed. Frontiers, Gif-sur-Yvette, France, 1991, 274 pp. (Basics of).
- [24] Y. Mehtar-Tani, C.A. Salgado, K. Tywoniuk, Phys. Rev. Lett. 106 (2011) 122002, arXiv:1009.2965 [hep-ph].
- [25] Y. Mehtar-Tani, C.A. Salgado, K. Tywoniuk, Phys. Lett. B 707 (2012) 156, arXiv:1102.4317 [hep-ph].
- [26] J. Casalderrey-Solana, E. Iancu, J. High Energy Phys. 1108 (2011) 015, arXiv:1105.1760 [hep-ph].
- [27] N. Armesto, H. Ma, Y. Mehtar-Tani, C.A. Salgado, K. Tywoniuk, J. High Energy Phys. 1201 (2012) 109, arXiv:1110.4343 [hep-ph].
- [28] Y. Mehtar-Tani, C.A. Salgado, K. Tywoniuk, J. High Energy Phys. 1204 (2012) 064, arXiv:1112.5031 [hep-ph].
- [29] Y. Mehtar-Tani, C.A. Salgado, K. Tywoniuk, J. High Energy Phys. 1210 (2012) 197, arXiv:1205.5739 [hep-ph].
- [30] Y. Mehtar-Tani, K. Tywoniuk, J. High Energy Phys. 1301 (2013) 031, arXiv:1105.1346 [hep-ph].
- [31] J. Casalderrey-Solana, Y. Mehtar-Tani, C.A. Salgado, K. Tywoniuk, Phys. Lett. B 725 (2013) 357, arXiv:1210.7765 [hep-ph].
- [32] J.-P. Blaizot, F. Dominguez, E. Iancu, Y. Mehtar-Tani, arXiv:1311.5823 [hep-ph].
- [33] Y.L. Dokshitzer, D.E. Kharzeev, Phys. Lett. B 519 (2001) 199.
- [34] N. Armesto, C.A. Salgado, U.A. Wiedemann, Phys. Rev. D 69 (2004) 114003.
- [35] B.W. Zhang, E. Wang, X.N. Wang, Phys. Rev. Lett. 93 (2004) 072301.
- [36] M. Djordjevic, M. Gyulassy, Nucl. Phys. A 733 (2004) 265.
- [37] B.I. Abelev, et al., STAR Collaboration, Phys. Rev. Lett. 98 (2007) 192301.
- [38] A. Adare, et al., PHENIX Collaboration, Phys. Rev. C 84 (2011) 044905, arXiv:1005.1627 [nucl-ex].
- [39] B. Abelev, et al., ALICE Collaboration, J. High Energy Phys. 1209 (2012) 112, arXiv:1203.2160 [nucl-ex].
- [40] B. Abelev, et al., ALICE Collaboration, Phys. Rev. Lett. 109 (2012) 112301, arXiv:1205.6443 [hep-ex].
- [41] S. Chatrchyan, et al., CMS Collaboration, J. High Energy Phys. 1205 (2012) 063, arXiv:1201.5069 [nucl-ex].
- [42] S. Chatrchyan, et al., CMS Collaboration, arXiv:1312.4198 [nucl-ex].