

ORIGINAL RESEARCH

Genetic identification of gastrointestinal parasites in the world's most endangered ungulate, the hirola (*Beatragus hunteri*)

Susana Remesar^{1,2}  | David García-Dios¹  | Giovanni Forcina^{3,4,5}  |
Abdullahi H. Ali^{6,7,8} | Mathew Ndunda⁹ | Michael J. Jowers^{3,4,10} 

¹Investigación en Sanidad Animal: Galicia (Grupo INVESAGA), Faculty of Veterinary Sciences, Universidade de Santiago de Compostela, Lugo, Spain

²IBADER—Instituto de Biodiversidade Agraria e Desenvolvemento Rural, Lugo, Spain

³CIBIO, Centro de Investigação em Biodiversidade e Recursos Genéticos, InBIO Laboratório Associado, Campus de Vairão, Universidade do Porto, Porto, Portugal

⁴BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, Universidade do Porto, Porto, Portugal

⁵Departamento de Ciencias de la Vida, Universidad de Alcalá, Global Change Ecology and Evolution Research Group (GloCEE)/Soil Biology and Subterranean Ecosystems Research Group (GIBSES), Alcalá de Henares, Madrid, Spain

⁶Department of Zoology and Physiology, University of Wyoming, Laramie, Wyoming, USA

⁷National Museums of Kenya, Nairobi, Kenya

⁸Hirola Conservation Programme, Garissa, Kenya

⁹Department of Veterinary and Capture Services, Kenya Wildlife Service, Nairobi, Kenya

¹⁰Departamento de Biología y Geología, Física y Química Inorgánica, Universidad Rey Juan Carlos, Madrid, Spain

Correspondence

Susana Remesar, Investigación en Sanidad Animal: Galicia (Grupo INVESAGA), Faculty of Veterinary Sciences, Universidade de Santiago de Compostela, Lugo, Spain.
Email: susana.remesar@usc.es

Funding information

Programme for Consolidating and Structuring Competitive Research Groups, Grant/Award Number: ED431C2023/16; Spanish Government Ministry of Universities; Portuguese Foundation for Science and Technology, Grant/Award Numbers: SFRH/BPD/109148/2015, PTDC/BAA-AGR/28866/2017

Abstract

Background: The hirola (*Beatragus hunteri*) is the world's most critically endangered antelope. Its parasitic fauna has never been genetically characterised, raising questions about its possible role in the species' population decline. This study aimed to assess the presence of gastrointestinal parasites in hirolas and their transmission pathways.

Methods: Thirty-one hirola faecal samples were examined using coprological methods. The identification of trematode and nematode species was performed by PCR amplification of the internal transcribed spacer 2 (ITS-2) and 18S ribosomal RNA (18S rRNA) genes.

Results: Trematode and nematode eggs were detected in five (16.1%) and 23 (74.2%) samples, respectively. The trematode sequence was identical to *Cotylophoron cotylophorum*, while the nematode sequence showed a 99.3% identity to *Chabaudstrongylus ninhae* (18S rRNA) and 99.4% identity to *Cooperia curticei* (ITS-2).

Limitations: The sample size is relatively small; however, it represents a large portion of the extant hirola population at the year of sampling.

Conclusions: A high percentage of the animals were infected by parasites whose presence may be associated with co-occurrence between hirolas and other ungulate species. Monitoring the parasite burden in local livestock may therefore be crucial for hirola conservation.

KEYWORDS

endangered species, gastrointestinal parasites, Paramphistomidae, Trichostrongylidae

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial](https://creativecommons.org/licenses/by-nc/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

© 2025 The Author(s). *Veterinary Record* published by John Wiley & Sons Ltd on behalf of British Veterinary Association.

INTRODUCTION

Gastrointestinal parasite infections in ungulates are economically and medically significant worldwide.¹ Among them, those caused by trematodes (Paramphistomidae and Fasciolidae), nematodes (i.e., Trichostrongylidae) and some protozoa are of most concern for both domestic and wild ungulates,² with several of these parasite species causing serious diseases,³ of which some are zoonotic.⁴

Some of these parasites have an indirect lifecycle where both intermediate and definitive hosts are necessary for the infection to occur. Consequently, all of the hosts play an important role in pathogen survival, proliferation and transmission, which are affected by several biotic and abiotic factors, including urban encroachment, global warming and faunal reshuffling.² Indeed, the free-living environmental stages of these parasites are significantly influenced by climate, with soil moisture being essential for the development and survival of the parasites as well as their intermediate and paratenic hosts. Therefore, while rising temperatures may accelerate the development of certain parasite larval stages,⁵ rainfall and vegetation could increase host exposure to parasites and affect their infection patterns.⁶ In addition, while some gastrointestinal parasites are host specific, others are generalists that can be transmitted between different hosts.^{7,8}

In recent years, parasitic infections have come to represent an emerging threat to several wild ungulates, posing a challenge to their management and ongoing conservation efforts.⁹ For instance, helminth parasitic burden is deemed to be one of the leading factors behind the drastic decline of some species,¹⁰ with bleak forecasts of climate warming-driven parasitic infections and associated disease outbreaks in certain regions.¹¹ In this context, characterising the gastrointestinal parasite communities in co-occurring livestock and wild ungulate species is crucial.¹²

The hirola (*Beatragus hunteri*) (Figure 1), first described in 1888,¹³ is a critically endangered medium-sized antelope with a narrow distribution currently restricted to northeastern Kenya but formerly encompassing bordering Somalia. This region is also inhabited by other ungulates, such as the oryx (*Oryx* spp.), the Grant's gazelle (*Nanger granti*), the Burchell's zebra (*Equus quagga burchellii*) and the coastal topi (*Damaliscus lunatus*). In addition, although hirolas avoid direct association with livestock, some individuals roam areas that have been previously grazed by cattle.¹⁴ To date, knowledge about hirola parasites is still limited; therefore, it is not known if these could have played a role behind its decline, which has been historically attributed to rinderpest virus transmission from other ruminants¹⁵ and, more recently, to habitat degradation and predation.^{16,17} Habitat restoration and protection are ongoing conservation efforts to enhance the recovery of the hirola, but characterising the diversity of parasite species infecting this antelope across its narrow range is crucial for understanding

potential health threats and enhancing conservation management.

For this reason, the main objective of the present study was to assess the presence of gastrointestinal parasites in the hirola and to molecularly identify them. The information obtained will help to elucidate infection pathways from other wild or domestic ungulates and facilitate the development of strategies to reduce this phenomenon.

MATERIALS AND METHODS

Study area and samples

A total of 39 hirola faecal samples from semi-captive herds within the predator-free fenced sanctuary in Garissa County (SANT; $n = 27$) and from wild herds at Ishaqbini Community Conservancy (CONS; $n = 5$) were collected from the ground between December 2017 and February 2018 (Figure 2). No metadata were recorded for seven samples. All the samples were individually labelled and preserved in plastic vials filled with 70% ethanol.

The faecal samples were genotyped to identify the individuals they came from, as described in a previous study.¹⁸ Eight of the 39 samples had non-unique genotypes, indicating that the same individuals had been sampled more than once, and these were therefore excluded from downstream analyses. The number of individuals identified per sampling area varied between 20 (SANT) and five (CONS), but for six individuals, we could associate no locality information. The final dataset consisted of faecal samples from 31 individuals, representing a large portion (around 25%) of the extant hirola population at the time of the survey.

Coprological techniques

Faecal samples were examined using two classical copromicroscopic methods: sedimentation^{20,21} and flotation.²² For both techniques, 3 g of faeces was homogenised in 42 mL of water and filtered using a 150 μm mesh. Two 12 mL crystal tubes were filled with this faecal solution, and the remaining solution was preserved in labelled plastic tubes stored at 4°C. These crystal tubes were then centrifuged at 680 \times g for 5 minutes and the supernatant was discarded.

Trematode eggs were detected by diluting the sediment contained in a single 12 mL crystal tube in 1000 mL of tap water. The suspension was allowed to settle for 20 minutes in a sedimentation cone; afterwards, the supernatant was discarded, and the sediment was resuspended in 500 mL of tap water. After another 20 minutes of sedimentation and the subsequent decantation, the solid phase was recovered into a 100 mL sedimentation cone that was filled with tap water. Subsequently, all the sediment obtained was used to fill a McMaster chamber, and this was directly observed under a microscope (Olympus CX43, Olympus Corporation). For the detection



FIGURE 1 Adult hirola (*Beatragus hunteri*)




FIGURE 2 Collection sites of hirola faecal samples. Modified from OpenStreetMap¹⁹

of nematode eggs, the sediment obtained from the other crystal tube was processed using flotation in Sheather's sucrose solution²² (specific gravity: 1.27) and observed under a microscope.

DNA extraction

Samples with excessive faecal material hindering egg collection were diluted and cleared with tap water. Gastrointestinal trematode and nematode eggs were individually collected from the faecal solution with a

micropipette and stored in RNase- and DNase-free plastic tubes at 4°C. DNA extraction was carried out in batches of four to five trematode or nematode eggs using the High Pure PCR Template Preparation Kit (Roche Diagnostics), following the manufacturer's instructions.

Parasite molecular detection

Trematode identification was performed using a previously described PCR protocol,²³ which targets an

TABLE 1 Primers used for Strongylidae species detection and identification

Gene target	Name	Primer sequence 5'–3'	Fragment size	Nt	T _m (°C)	CG (%)	Self-dimers	Cross-primer dimers
18S rRNA	StrongF	AAGGGTGCAATTATTAGAGCA	650 bp	21	60.6	38.1	–	–
	StrongR	GAACCGAAGTCCTCTTTTATTAT		23	58.7	34.8	–	–

Abbreviations: CG, cytosine–guanine content; Nt, number of nucleotides; T_m, melting temperature.

approximately 400 bp-long fragment of the internal transcribed spacer 2 (ITS-2) gene. In each amplification reaction, nuclease-free water was included as a negative control, and rumen fluke (*Calicophoron daubneyi*) DNA obtained from a cow (*Bos taurus*) was included as a positive control.

For identifying gastrointestinal nematode parasites, a new primer set (StrongF/StrongR) was designed to target approximately 650 bp of the 18S ribosomal RNA (18S rRNA) gene of the species belonging to the family Strongylidae. Nucleotide sequence variation across different genera, as inferred from their available GenBank records (Table S1), was used for designing the primers after aligning the sequences with the online CLUSTALW tool GenomeNet Multiple Sequence Alignment (www.genome.jp/tools-bin/clustalw). Primer compatibility and melting temperatures were assessed using the online tool Multiple Primer Analyser (ThermoFisher Scientific) (Table 1), while Primer-Basic Local Alignment Search Tool (BLAST, www.ncbi.nlm.nih.gov/tools/primer-blast/) was used to assess their target sequence identity and amplifiability. The PCR reaction mix consisted of 1.75 µL of 3.5 mM MgCl₂, 2.5 µL of 200 µM dNTPs, 1.5 µL (0.6 µM) of each primer, 0.5 units of NZYtaq II DNA polymerase (NZYTech) and 3 µL of template DNA in a final volume of 25 µL. Amplifications were carried out in a T100 Thermal Cycler (Bio-Rad Laboratories). The cycling conditions were as follows: initial denaturation of 5 minutes at 94°C, 40 cycles of 45 seconds at 94°C, 45 seconds at 59°C and 1 minute at 72°C followed by a final elongation of 7 minutes at 72°C. For further identification of the gastrointestinal nematodes, a PCR protocol previously described,²⁴ targeting an approximately 300 bp fragment of the ITS-2 gene, was carried out. In each amplification reaction, nuclease-free water was included as a negative control, and *Trichostrongylus colubriformis* DNA obtained from a sheep (*Ovis aries*) was included as a positive control.

Successful amplification was assessed using electrophoresis on 1.5% agarose gels stained with RedSafe (iNtRON Biotechnology) and then visualised using a Fluor-S MultiImager (Bio-Rad Laboratories). PCR products were subsequently purified and sequenced bidirectionally at the Sequencing and Fragment Analysis Unit of Santiago de Compostela University. The sequences were manually edited using ChromasPro (Technelysium) and compared with those deposited in the GenBank database using BLAST (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>). The unique partial sequences identified in this study were deposited in GenBank

under accession numbers PQ639947, PQ639948 and PQ640031.

RESULTS

A total of 23 individuals (74.2%; 95% confidence interval [CI]: 55.1–87.5%) tested parasite positive using at least one coprological technique. Specifically, trematode eggs were detected through sedimentation in five of the 31 individuals (16.1%; 95% CI: 6.1–34.5%), while gastrointestinal nematode eggs were detected through sucrose flotation in 23 individuals (74.2%; 95% CI: 55.1–87.5%) (Figure 3). No other parasitic forms were detected in the samples analysed. Coinfections with nematodes were observed in all of five the animals positive for trematodes. The overall prevalence of gastrointestinal parasites varied across localities, with CONS showing the highest prevalence (5/5; 100%; 95% CI: 46.3–100%), followed by SANT (16/20; 80%; 95% CI: 55.7–93.4%) and individuals of uncertain provenance (2/6; 33.3%; 95% CI: 6.0–75.9%). The gastrointestinal nematode prevalence was the same as the total prevalence for each sampling area, being 100%, 80% and 33.3% in individuals sampled in CONS, SANT and uncertain provenance, respectively. Meanwhile, trematode prevalence was similar in CONS (1/5; 20%; 95% CI: 1.1–70.1%) and SANT (4/20; 20%; 95% CI: 6.6–44.3%), but no trematodes were detected in individuals of uncertain provenance.

The Paramphistomidae trematode ITS-2 sequence (381 bp) obtained was 100% identical to *Cotylophoron cotylophorum* (GenBank record KX668959) isolated in cattle from Kenya²⁵ and showed a high similarity (99.5–99.7%) to *Cotylophoron* sp. (GenBank records KX668913, KX668970 and KX668975) isolated in cattle and sheep from Kenya.²⁵

Although a few nematode eggs were collected and five DNA extractions were carried out, only one sample tested positive using the PCR protocol targeting the Strongylidae 18S rRNA gene. This sample showed 99.3% identity to *Chabaudstrongylus ninhae*, syn. *Cooperia ninhae* (GenBank record LC415111), which was isolated from a Reeves's muntjac (*Muntiacus reevesi*) in Japan.²⁶ This newly amplified sequence also showed a 98.4% similarity to *Trichostrongylus colubriformis* (GenBank record AJ920350) obtained from sheep (*O. aries*)²⁷ and a 98.0% similarity to *Haemonchus contortus* (GenBank records EU086374 and EU086375) obtained from a captive giraffe (*Giraffa camelopardalis*) from Florida,²⁸ an *H. contortus* laboratory strain (GenBank records LS997563




FIGURE 3 Examples of (a) trematode and (b) nematode eggs observed in the faecal samples

and LS997564)²⁹ and *Graphidioides affinis* (GenBank records OR611714 and OR611716) obtained from a Patagonian mara (*Dolichotis patagonum*) from Bulgaria (unpublished). The ITS-2 sequence (325 bp) obtained for this sample showed a 98.2–99.4% identity to the previously deposited *Cooperia curticei* sequences (GenBank records KC998734–KC998739) isolated in sheep from New Zealand,³⁰ and a 97.2% identity to *Cooperia oncophora* sequences (GenBank records KC998742, KC998743 and AB534601) obtained from cattle in New Zealand³⁰ and Japan (unpublished). This sequence also showed a 96.6% similarity to the *C. ninhae* sequence (GenBank record LC415111) isolated from a Reeves's muntjac in Japan.²⁶

DISCUSSION

The hirola—listed as critically endangered in the IUCN Red List of Threatened Species and still experiencing a decreasing demographic trend³¹—is referred to as the most threatened antelope in the world. Its dramatic global population decline has been attributed to multiple causes, among which the 1980s rinderpest virus outbreak, resulting in mass mortality of the ruminants in eastern Kenya,³² stands out. While the recovery of the hirola started only after the eradication of this virus, which points to its leading role in the plight faced by this rare antelope, several other factors, such as habitat loss and degradation, poaching and competition with livestock, have also been involved.^{17,33,34} Although some authors have suggested that the limited recovery of hirola global populations is likely attributable to their low genetic diversity,¹⁵ the only pertinent molecular investigation carried out until now revealed no evidence of inbreeding.¹⁸

The present study is, to the authors' knowledge, the first to assess the presence of gastrointestinal parasites in the hirola through sequencing, thus advancing our knowledge on the ecology of this rare species. The results obtained indicate that more than half of the studied animals were infected with gastrointestinal parasites. These findings agree with those reported for other wild ungulates from Kenya, where more than 50% of the surveyed individuals carried at

least one helminth species,³⁵ suggesting a high exposure to these parasites. Prevalences up to 100% were reported in Burchell's zebras^{35,36} and several African antelopes,^{35,37,38} including hirolas, while lower, but still high, values were reported in zebuine cattle (*Bos indicus*) (29%) and Cape buffalo (*Syncerus caffer*) (47–87%).^{35,37} On the other hand, helminths were only found in 9% of the examined giraffes.³⁵

Two taxa of gastrointestinal parasites were detected by coprological analysis of the samples collected for this study: Paramphistomidae trematodes and Trichostrongylidae nematodes. Previous studies in hirolas have reported gastrointestinal protozoa (Coccidia) together with Trichostrongylidae, Strongylidae and Ascarididae species, bronchopulmonary nematodes and trematodes.^{38,39} However, other nematodes (Trichuridae, Spiruridae, Oxyuridae) and cestodes (Anoplocephalidae) detected in other wild ungulates from Kenya^{35,38–40} have not been identified in this species. We cannot rule out that the low parasite diversity seen in the present study could be due to the sample preservation method rather than reflecting an intrinsically poor gastrointestinal parasite community. Nevertheless, low parasite diversity has been reported in other wild ungulates from the same area.^{37,40} According to previous observations,⁴¹ this outcome may be due to the animals' limited contact with different habitats and parasite hosts, which is in line with the narrow distribution range of the hirola. Concordantly, it has been proposed that migration and transhumance could lead to higher parasite diversity³⁵; therefore, the low parasite diversity observed in this species could also be associated with absence of seasonal movements of the sampled herds at the time of sample collection.

Gastrointestinal nematodes (74.2%) were more frequent than trematodes (16.1%), which is consistent with previous studies carried out on hirolas, which found prevalences of around 54% and 24% for these taxa, respectively.^{38,39} These results are also consistent with those obtained for other African antelopes^{35,37,40–42} and Cape buffaloes.³⁷

As far as the high prevalence of Trichostrongylidae nematodes is concerned, it is important to note their ability to survive in dry conditions.^{43,44} In addition,

it has been reported that some ungulates share certain gastrointestinal nematodes species,^{35,45,46} which increases egg environmental contamination and the probability of interspecies transfer. By contrast, Paramphistomidae trematodes require wet environments to survive, since they need an amphibious or aquatic gastropod to complete their lifecycle.⁴⁴ These parasites have been detected in a high percentage of water buffaloes (*Bubalus bubalis*),^{35,37} which inhabit marshy areas where Paramphistomidae intermediate hosts⁴⁷ thrive. In addition, species of this family have also been detected in up to 90% of the domestic cattle, sheep and goats from Kenya,^{25,40} which inhabit watered natural habitats where the intermediate host is present.²⁵

Although trematode eggs had been previously detected in hirola faecal samples,³⁸ no species identification had been carried out. It is worth mentioning that *Cotylophoron* spp. had been previously detected in a wide range of domestic and wild African ruminants.^{25,48,49} Given that *C. cotylophorum* is known for its broad host range and is frequently found in African ungulates,⁴⁹ its identification in the hirola is not unexpected, especially given the co-occurrence of this antelope with other wild and domestic ruminants,^{14,50} which could facilitate cross-infection between species. This is in line with previous studies, which suggested that the infection risk with broad host-range parasites increases in communities with co-occurring hosts.^{37,46} This cross-species contact is of particular concern for endangered species that are the object of intensive conservation efforts.³⁷ However, previous studies have suggested that deworming domestic species could be an effective way to reduce the transmission between livestock and threatened wild ungulates.^{46,51}

The characterisation of the 18S rRNA gene revealed that the Trichostrongylidae species detected in the present study was phylogenetically close to a *C. ninhiae* specimen obtained from a Reeves's muntjac from Japan.²⁶ The genus *Chabaudstrongylus* has been previously identified in some deer (Cervidae) and mouse deer (Tragulidae) species,^{52–54} suggesting that the eggs found in the hirola faecal samples might therefore belong to the same or a highly related genus. However, the characterisation of the partial sequence obtained for the ITS-2 gene found the highest similarity to be with *C. curticei*, a species commonly detected in sheep and goats,⁴⁴ with a sequence identity of lower than 97% for the closest *C. ninhiae* sequence.

Since the hirola is the only representative of the genus *Beatragus*,^{55,56} the detected nematode could be host specific; however, information about the molecular characterisation of Strongylidae species in African antelopes is still scarce. A previous study carried out in the hirola³⁸ identified other Trichostrongylidae genera, such as *Trichostrongylus* spp., *Oesophagostomum* spp., *Bunostomum* spp., *Haemonchus* spp., *Cooperia* spp. and *Ostertagia* spp.; however, there is no information about the procedures used for such identification, more so as nematode eggs of those gen-

era are morphologically indistinguishable. In view of these considerations and the limited number of eggs that could be collected and analysed in this study, it is conceivable that other Trichostrongylidae species could be co-circulating in the same hirola population. Moreover, we cannot rule out that the partial sequences obtained belong to two different species. Several *Oesophagostomum*, *Trichostrongylus*, *Cooperia*, *Haemonchus* and *Teladorsagia* species have been molecularly identified in African ruminants,^{40,57,58} which calls for further studies to assess the systematic placement of hirola Trichostrongylidae as well as to unravel their epidemiology and host range.

Finally, it is important to state that the samples processed in the present investigation were retrieved from the environment and preserved with the purpose of carrying out a genetic diversity study of the extant hirola populations, not to perform coprological analysis. Ethanol preservation is a good method for conserving faecal samples over time if DNA retrieval is the main objective,⁵⁹ but the best protocol for studies such as this one is to refrigerate and process them as soon as possible to prevent the death, destruction and morphological alteration of the parasitic forms.^{60,61} For this reason, the presence of bronchopulmonary nematodes was not assessed in the present investigation, since their detection technique relies on the existence of motile larvae.⁶² In addition, although sedimentation is a quantitative technique, it was only used in a qualitative way, since the preservation method could have affected parasite counts, resulting in an underestimation of the intensity of egg shedding. Nevertheless, we deem our results to still hold a special interest from an ecological point of view, since data on gastrointestinal helminths affecting the hirola are still scant.

In conclusion, our study highlights the presence of two gastrointestinal helminth groups, Paramphistomidae and Trichostrongylidae, in the critically endangered hirola antelope. These findings underscore the need for further research to understand the pathogenicity and epidemiology of these helminths in hirola populations, which is essential for effective conservation.

AUTHOR CONTRIBUTIONS

Susana Remesar designed and carried out the analyses and wrote the manuscript. David García-Dios carried out analyses and reviewed the manuscript. Giovanni Forcina performed data curation and writing—review and editing. Abdullahi H. Ali and Mathew Ndunda performed sampling and writing—review and editing. Michael J. Jowers conceived the study and reviewed and edited the manuscript.

ACKNOWLEDGEMENTS

We are grateful to the Kenya Wildlife Service (KWS) and the Hirola Conservation Programme for permit issuance, and to Francis Gakuya (KWS Head of Veterinary Services), Moses Otiende and Isaac Lekoolool for assisting in sample collection and shipment. This

research was partially supported by the Programme for Consolidating and Structuring Competitive Research Groups (ED431C2023/16; Xunta de Galicia, Spain), the Spanish Government Ministry of Universities (“María Zambrano”—Next Generation EU grants to G.F. and M.J.J.) and by fellowships from the Portuguese Foundation for Science and Technology awarded to M.J.J. (FCT, SFRH/BPD/109148/2015) and G.F. (FCT, PTDC/BAA-AGR/28866/2017).

CONFLICT OF INTEREST STATEMENT

The authors declare they have no conflicts of interest.


DATA AVAILABILITY STATEMENT

The authors will make the data available upon request.

ETHICS STATEMENT


This study does not involve animal experimentation, so no ethical approval was deemed necessary.

ORCID

Susana Remesar  <https://orcid.org/0000-0002-8071-3806>

David García-Dios  <https://orcid.org/0000-0001-6364-766X>

Giovanni Forcina  <https://orcid.org/0000-0001-5727-7770>

Michael J. Jowers  <https://orcid.org/0000-0001-8935-5913>

REFERENCES

- Waller PJ. From discovery to development: current industry perspectives for the development of novel methods of helminth control in livestock. *Vet Parasitol.* 2006;139(1–3):1–14.
- Bowman DD. *Georgis' parasitology for veterinarians*. 10th ed. Philadelphia: Saunders; 2016.
- Fowler ME, Bravo PW. *Parasites. Medicine and surgery of camelids*. 3rd ed. Ames, Iowa: Wiley-Blackwell; 2010.
- Nyagura I, Malatji MP, Mukaratirwa S. Occurrence of *Fasciola* (Digenea: Fasciolidae) species in livestock, wildlife and humans, and the geographical distribution of their intermediate hosts in South Africa—a scoping review. *Front Vet Sci.* 2022;9:935428.
- Rose H, Wang T, van Dijk J, Morgan ER. GLOWORM-FL: a simulation model of the effects of climate and climate change on the free-living stages of gastrointestinal nematode parasites of ruminants. *Ecol Modell.* 2015;297:232–45.
- Shearer CL, Ezenwa VO. Rainfall as a driver of seasonality in parasitism. *Int J Parasitol Parasites Wildl.* 2020;12:8–12.
- Arias-Pacheco C, Pezo D, Mathias LA, Tebaldi JH, Castelo-Oviedo H, Lux-Hoppe EG. Parasitological status of vicuñas (*Vicugna vicugna*) from southeastern Peru and its relationship with fiber quality. *Trop Anim Health Prod.* 2021;53(2):211.
- Cañal V, Beltrame MO. Gastrointestinal parasite diversity of South American camelids (Artiodactyla: Camelidae): first review throughout the native range of distribution. *Int J Parasitol Parasites Wildl.* 2022;19:222–42.
- Daszak P, Cunningham AA, Hyatt AD. Emerging infectious diseases of wildlife—threats to biodiversity and human health. *Science.* 2000;287(5452):443–49.
- Morgan ER, Shaw SE, Brennan SF, De Waal TD, Jones BR, Mulcahy G. *Angiostrongylus vasorum*: a real heart breaker. *Trends Parasitol.* 2006;21(2):49–51.
- Kutz SJ, Hoberg EP, Nagy J, Polley L, Elkin B. “Emerging” parasitic infections in arctic ungulates. *Integr Comp Biol.* 2004;44(2):109–18.
- Bhat RA, Tak H, Bhat BA, Dar JA, Ahmad R. Gastrointestinal helminth parasites of wild ungulates in Hirpora Wildlife Sanctuary, Kashmir, India. *J Parasit Dis.* 2022;46(3):804–10.
- Slater PL. Description of Hunter's antelope. *Proc Zool Soc.* 1889;58:372–77.
- Bunderson WT. *Ecological separation of wild and domestic mammals in an East African Ecosystem*. Logan: Utah State University; 1981.
- Butynski TM. Independent evaluation of hirola antelope (*Beatragus hunteri*) conservation status and conservation action in Kenya. Nairobi: Kenya Wildlife Service and Hirola Management Committee; 2000.
- Ali AH, Ford AT, Evans JS, Mallon DP, Hayes MM, King J, et al. Resource selection and landscape change reveal mechanisms suppressing population recovery for the world's most endangered antelope. *J Appl Ecol.* 2017;54(6):1720–29.
- Ali AH, Kauffman MJ, Amin R, Kibara A, King J, Mallon D, et al. Demographic drivers of a refugee species: large-scale experiments guide strategies for reintroductions of hirola. *Ecol Appl.* 2018;28(2):275–83.
- Jowers MJ, Queirós J, Pinto RR, Ali AH, Ndunda M, Angelone S, et al. Genetic diversity in natural range remnants of the critically endangered hirola antelope. *Zool J Linn Soc.* 2020;190:384–95.
- OpenStreetMap Contributors. Planet dump. 2022. Available from: <https://www.openstreetmap.org>
- Rehbein S, Kokott S, Lindner, T. Evaluation of techniques for the enumeration of *Dicrocoelium* eggs in sheep faeces. *Zentralbl Veterinarmed.* 1999;46(3):133–40.
- García-Dios D, Panadero R, Díaz P, Viña M, Remesar S, Prieto A, et al. The goat as a risk factor for parasitic infections in ovine flocks. *Animals.* 2021;11(7):2077.
- Foreyt W. *Veterinary parasitology*. Iowa: Blackwell Publishing; 1989.
- García-Dios D, Díaz P, Remesar S, Viña M, Martínez-Calabuig N, Saldaña A, et al. Prevalence, risk factors and molecular identification of paramphistomid species in sheep from a Spanish endemic area. *Ir Vet J.* 2024;77(1):21.
- Gasser RB, Chilton NB, Hoste H, Beveridge I. Rapid sequencing of rDNA from single worms and eggs of parasitic helminths. *Nucleic Acids Res.* 1993;21(10):2525–26.
- Laidemitt MR, Zawadzki ET, Brant SV, Mutuku MW, Mkoji GM, Loker ES. Loads of trematodes: discovering hidden diversity of paramphistomoids in Kenyan ruminants. *Parasitology.* 2017;144(2):131–47.
- Setsuda A, Kato E, Sakaguchi S, Tamemasa S, Ozawa S, Sato H. *Chabaudstrongylus ninhiae* (Trichostrongylidae: Cooperiinae) and *Oesophagostomum muntiacum* (Chabertiidae: Oesophagostominae) in feral alien Reeves's muntjacs on Izu-Oshima Island, Tokyo, Japan. *J Helminthol.* 2019;94:e48.
- Chilton NB, Huby-Chilton F, Gasser RB, Beveridge I. The evolutionary origins of nematodes within the order Strongylida are related to predilection sites within hosts. *Mol Phylogenet Evol.* 2006;40(1):118–28.
- Garretson PD, Hammond EE, Craig TM, Holman PJ. Anthelmintic resistant *Haemonchus contortus* in a giraffe (*Giraffa camelopardalis*) in Florida. *J Zoo Wildl Med.* 2009;40(1):131–39.
- Laing R, Kikuchi T, Martinelli A, Tsai IJ, Beech RN, Redman E, et al. The genome and transcriptome of *Haemonchus contortus*, a key model parasite for drug and vaccine discovery. *Genome Biol.* 2013;14(8):R88.
- Bisset SA, Knight JS, Bouchet CL. A multiplex PCR-based method to identify strongylid parasite larvae recovered from ovine faecal cultures and/or pasture samples. *Vet Parasitol.* 2014;200(1–2):117–27.
- IUCN SSC Antelope Specialist Group. *Beatragus hunteri*. The IUCN Red List of Threatened Species 2017: e.T6234A50185297. 2017. Available from: <https://www.iucnredlist.org/species/6234/50185297>. Accessed on 9 Dec 2019.
- Kock ND, Kock RA, Wambua J, Mwanzia J. Pathological changes in free-ranging African ungulates during a rinderpest epizootic in Kenya, 1993 to 1997. *Vet Rec.* 1999;145(18):527–28.

33. Andanje SA. Factors limiting the abundance and distribution of hirola (*Beatragus hunteri*) in Kenya. Unpublished Ph.D. Thesis. Newcastle University; 2002.
34. Ali AH, Goheen JR, Evans JS, Hayes MM, Kibara A, Amin R. Hirola conservation project progress report: primary findings from 2012–2014. 2014. Available from: <https://ptes.org/wp-content/uploads/2015/02/Kenya-hirola-progress-report-2012-2014.pdf>
35. VanderWaal K, Omondi GP, Obanda V. Mixed-host aggregations and helminth parasite sharing in an East African wildlife-livestock system. *Vet Parasitol.* 2014;205(1–2):224–32.
36. Wambwa E, Ogara WO, Mudakha D. A comparative study of gastrointestinal parasites between ranched and free ranging Burchell's zebra (*Equus burchelli antiquorum*) in Isiolo district, Kenya. *J Vet Sci.* 2004;5(3):215–20.
37. Ezenwa VO. Habitat overlap and gastrointestinal parasitism in sympatric African bovids. *Parasitology.* 2003;126(4):379–88.
38. Njeru JM, Kariuki E, Andanje SA, Wahungu GM, Wanjala PM, Wahungu GM. Prevalence and intensity of gastrointestinal parasites Hirola (*Beatragus hunteri*) and livestock in Southern Kenya. *Afr J Educ Sci Technol.* 2014;1:175–84.
39. Njeru JM, Mutevi WP, Wahungu GM, Andanje SA. Spatial and temporal variations in gastrointestinal parasitism in the critically endangered hirola and livestock in Southern Kenya. *J Nat Sci Res.* 2016;6:85–93.
40. Obanda V, Maingi N, Muchemi G, Ng'ang'a CJ, Angelone S, Archie EA. Infection dynamics of gastrointestinal helminths in sympatric non-human primates, livestock and wild ruminants in Kenya. *PLoS One.* 2019;14(6):e0217929.
41. Bordes F, Morand S. Parasite diversity: an overlooked metric of parasite pressures? *Oikos.* 2009;118(6):801–6.
42. Kariuki DP, Injairo R, Boyce WL, Welde BT, Ngethe S. Parasite survey of 8 wild animals in the Ruma-National Park. *Ann Trop Med Parasitol.* 1989;83(suppl. 1):115–18.
43. Andersen FL, Levine ND. Effect of desiccation on survival of free-living stages of *Trichostrongylus colubriformis*. *J Parasitol.* 1968;54(1):117–28.
44. Deplazes P, Eckert J, Mathis A, von Samson-Himmelstjerna G, Zahner H. Parasitology in veterinary medicine. Netherlands: Wageningen Academic Publisher; 2016
45. Mijele D, Iwaki T, Chiyo PI, Otiende M, Obanda V, Rossi L, et al. Influence of massive and long distance migration on parasite epidemiology: lessons from the great wildebeest migration. *Ecohealth.* 2016;13(4):708–19.
46. Titcomb GC, Pansu J, Hutchinson MC, Tombak KJ, Hansen CB, Baker CCM, et al. Large-herbivore nemabiomes: patterns of parasite diversity and sharing. *Proc Biol Sci.* 2022;289(1974):20212702.
47. Penzhorn BL. Coccidian oocyst and nematode egg counts of free-ranging African buffalo (*Syncerus caffer*) in the Kruger National Park, South Africa. *J S Afr Vet Assoc.* 2000;71(2):106–8.
48. Keyyu JD, Kassuku AA, Msalilwa LP, Monrad J, Kyvsgaard NC. Cross-sectional prevalence of helminth infections in cattle on traditional, small-scale and large-scale dairy farms in Iringa district, Tanzania. *Vet Res Commun.* 2006;30(1):45–55.
49. Pfukenyi DM, Mukaratirwa S. Amphistome infections in domestic and wild ruminants in East and Southern Africa: a review. *Onderstepoort J Vet Res.* 2018;85(1):e1–e13.
50. Andanje SA, Ottichilo WK. Population status and feeding habits of the translocated sub-population of Hunter's antelope or hirola (*Beatragus hunteri*, Sclater, 1889) in Tsavo East National Park, Kenya. *Afr J Ecol.* 1999;37(1):38–48.
51. Titcomb G, Hulke J, Mantas JN, Gituku B, Young H. Cattle aggregations at shared resources create potential parasite exposure hotspots for wildlife. *Proc Biol Sci.* 2023;290(2023):20232239.
52. Durette-Desset MC, Krishnasamy M. *Cooperia caballeroi* n. sp. Nematoda Trichostrongylidae, parasite of *Tragulus javanicus* en Malaisie. In: *Excerta Parasitologica en Memoria del Dr. Eduardo Caballero y Caballero.* Instituto de Biología de Mexico. Mexico. 1977. pp. 429–32.
53. Drózd J. *Cooperia ninhae* sp.n. (Nematoda: Trichostrongylidae), a parasite of *Muntjac muntjak* in Vietnam. *Acta Parasitol Pol.* 1967;14:209–11.
54. Inoue K, Shishida K, Kawarai S, Takeda S, Minami M, Taira K. Helminthes detected from wild sika deer (*Cervus nippon centralis*) in Kanto-Chubu region, Japan. *Parasitol Int.* 2021;87:102485.
55. Pitra C, Kock RA, Hofmann RR, Lieckfeldt D. Molecular phylogeny of the critically endangered Hunter's antelope (*Beatragus hunteri* Sclater, 1889). *J Zool Syst Evol Res.* 1997;36(4):179–84.
56. Steiner CC, Charter SJ, Houck ML, Ryder OA. Molecular phylogeny and chromosomal evolution of Alcelaphini (Antilopinae). *J Hered.* 2014;105(3):324–33.
57. Troell K, Mattsson JG, Alderborn A, Höglund J. Pyrosequencing analysis identifies discrete populations of *Haemonchus contortus* from small ruminants. *Int J Parasitol.* 2003;33(7):765–71.
58. Paguem A, Abanda B, Ngwasiri NN, Eisenbarth A, Renz A, Streit A, et al. Host specificity and phylogeny of Trichostrongylidae of domestic ruminants in the Guinea savannah of the Adamawa plateau in Cameroon. *Vet Parasitol Reg Stud Rep.* 2020;21:100412.
59. Blekhman R, Tang K, Archie EA, Barreiro LB, Johnson ZP, Wilson ME, et al. Common methods for fecal sample storage in field studies yield consistent signatures of individual identity in microbiome sequencing data. *Sci Rep.* 2016;6:31519.
60. Jagla E, Spiewak J, Zalesny G, Popiolek M. Effect of storage and preservation of horse faecal samples on the detectability and viability of strongylid nematode eggs and larvae. *Bull Vet Inst Pulawy.* 2013;57:161–65.
61. González M, de Ybáñez RR, Rodríguez-Caro RC, Maíz-García A, Gómez L, Giménez A, et al. Assessing DESS solution for the long-term preservation of nematodes from faecal samples. *Res Vet Sci.* 2022;153:45–48.
62. Ministry of Agriculture, Fisheries and Food. Manual of veterinary parasitological laboratory techniques. London: HMSO; 1986.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Remesar S, García-Dios D, Forcina G, Ali AH, Ndunda M, Jowers MJ. Genetic identification of gastrointestinal parasites in the world's most endangered ungulate, the hirola (*Beatragus hunteri*). *Vet Rec.* 2025;e5223. <https://doi.org/10.1002/vetr.5223>