

ORIGINAL ARTICLE

Learning to resuscitate at school. Study in 8-12 year-old schoolchildren[☆]



Santiago Martínez-Isasi^{a,b,c}, Cristian Abelairas-Gómez^{a,b,d}, María Pichel-López^e, Roberto Barcala-Furelos^{b,e}, Cristina Varela-Casal^{e,*}, David Vázquez-Santamariña^f, Luis Sánchez-Santos^g, Antonio Rodríguez-Núñez^{a,b,c,h}

^a Grupo de investigación CLINURSID, Departamento de Psiquiatría, Radiología y Salud Pública, Enfermería y Medicina, Universidade de Santiago de Compostela, Santiago de Compostela, Spain

^b Grupo de Soporte Vital y Simulación Médica, Instituto de Investigación Sanitaria de Santiago de Compostela (IDIS), Santiago de Compostela, Spain

^c Facultad de Enfermería, Universidade de Santiago de Compostela, Santiago de Compostela, Spain

^d Facultad de Ciencias de la Educación, Universidade de Santiago de Compostela, Santiago de Compostela, Spain

^e Grupo de Investigación REMOSS, Facultad de Ciencias de la Educación y del Deporte, Universidade de Vigo, Pontevedra, Spain

^f Servicio de Urgencias, Instituto de Investigación Biomédica de A Coruña (INIBIC), Complejo Hospitalario Universitario de A Coruña (CHUAC), Universidade da Coruña, A Coruña, Spain

^g Área Sanitaria Santiago de Compostela-Barbanza, Spain

^h Unidad de Cuidados Intensivos de Pediatría, Hospital Clínico Universitario de Santiago de Compostela, Universidade de Santiago de Compostela, Santiago de Compostela, Spain

Received 14 July 2020; accepted 27 September 2020

Available online 20 December 2021

KEYWORDS

Basic life support;
Cardiopulmonary
resuscitation;
Learning;
Schoolchildren;
Quality of chest
compression

Abstract

Objective: To quantitatively assess the learning capacity of school children aged between 8-12 years in basic life support (theory and practice) after a feasible school training programme.

Material and methods: Quasi-experimental study with a convenience sample of 567 pupils in 3rd and 5th year of Primary Education, and first year of Compulsory Secondary Education, from 3 public schools in Galicia. They received 2h (one theoretical and another practical) of basic life support training by their Physical Education teachers, as part of the school program. The children were evaluated by a theoretical test and a practical skill test that measured the quality of chest compressions, and assessed the performance of the basic life support sequence.

Results: The level of knowledge increased with respect to the baseline, and was higher in the higher grades ($P < .001$). The complete basic life support sequence was carried out by 16.5% of pupils in the 3rd year of Primary Education, 54.4% of pupils in the 5th year of Primary

[☆] Please cite this article as: Martínez-Isasi S, Abelairas-Gómez C, Pichel-López M, Barcala-Furelos R, Varela-Casal C, Vázquez-Santamariña D, et al. Aprendiendo a reanimar en la escuela. Estudio en escolares de 8-12 años. An Pediatr (Barc). 2022;96:17–24.

* Corresponding author.

E-mail address: cristinavarelacasal@uvigo.es (C. Varela-Casal).

PALABRAS CLAVE

Soporte vital básico;
Reanimación
cardiopulmonar
básica;
Enseñanza;
Escolares;
Calidad de las
compresiones
torácicas

Education, and 28.5% of pupils in the 1st year of Secondary Education ($P = .030$). The following compression quality parameters improved significantly with age: continuity of compressions ($P < .001$), percentage of compressions performed at correct depth ($P = .002$), and median depth ($P < .001$), while the percentage of compressions with correct decompression decreased significantly ($P < .001$).

Conclusions: Although their anthropometric characteristics may not allow them to achieve the ideal quality of this manoeuvre, a 2h theoretical and practical training programme, taught by Physical Education teachers, helps to improve the ability of children younger than 13 years old to recognise the emergency, start the chain of survival, and initiate chest compressions.

© 2021 Published by Elsevier España, S.L.U. on behalf of Asociación Española de Pediatría. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Aprendiendo a reanimar en la escuela. Estudio en escolares de 8-12 años

Resumen

Objetivo: Valorar cuantitativamente la capacidad de aprendizaje en soporte vital básico (teórica y práctica) de escolares de 8-12 años con un programa de formación adaptado a las escuelas.

Material y métodos: Estudio cuasiexperimental con una muestra de conveniencia de 567 alumnos de 3º y 5º de Educación Primaria y 1º de Educación Secundaria Obligatoria, de 3 colegios concertados de Galicia, que recibieron 2 h (una teórica y otra práctica) de formación en soporte vital básico por parte de sus profesores de Educación Física integrada en el programa escolar. Los niños fueron evaluados mediante un test teórico y una prueba práctica que midió la calidad de las compresiones torácicas y valoró la secuencia de soporte vital básico.

Resultados: El nivel de conocimientos aumentó respecto al basal y fue mayor en los alumnos de cursos superiores ($p < 0,001$). La secuencia completa de soporte vital básico fue realizada por el 16,5% de los escolares de 3º de Educación Primaria, el 54,4% de los de 5º de Educación Primaria y el 28,5% de los de 1º de Educación Secundaria Obligatoria ($p = 0,030$). Los siguientes parámetros de calidad de las compresiones mejoraron de forma significativa con la edad: continuidad de las compresiones ($p < 0,001$), porcentaje de compresiones realizadas con una profundidad correcta ($p = 0,002$) y mediana de profundidad ($p < 0,001$), mientras que el porcentaje de compresiones con descompresión correcta descendió significativamente ($p < 0,001$).

Conclusiones: Un programa formativo teórico-práctico de 2 h, impartido por los profesores de Educación Física, contribuye a mejorar la capacidad de los niños menores de 13 años para reconocer la emergencia, poner en marcha la cadena de supervivencia e iniciar las compresiones torácicas, aunque posiblemente sus características antropométricas no les permitan alcanzar la calidad ideal establecida para esta maniobra.

© 2021 Publicado por Elsevier España, S.L.U. en nombre de Asociación Española de Pediatría. Este es un artículo Open Access bajo la licencia CC BY-NC-ND (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Out-of-hospital cardiac arrest (OHCA) is witnessed by relatives, friends or other people in 70% of cases. In most instances, bystander cardiopulmonary resuscitation (CPR) attempts take place in the home, but CPR is only attempted before the arrival of medical emergency teams in less than 20% of cases,¹ when early CPR is a key determinant of survival and neurologic outcome.² The main strategy to increase the frequency of bystander CPR is training in basic life support (BLS), and in this regard, school-based training of students may be one of the most effective approaches.^{1,3}

In Spain, there have been several official initiatives and projects to provide BLS training in the school setting,⁴⁻⁶ but there is no evidence yet on its actual impact on bystander

initiation of BLS and victim survival. On the other hand, several studies in recent years have analysed the methodology of BLS training in children, with educational strategies varying significantly in terms of the duration of training, the methods used, the instructors and the minimum age for training.⁷

The *Kids Save Lives* initiative promoted by the European Resuscitation Council (ERC) with the support of the World Health Organization recommends 2 hours of school-based CPR training a year starting from age 12 years.³ However, it may not be necessary to wait until this age to start teaching first aid and BLS skills to children, as children as young as 6 years of age can learn the chain of survival⁸ and even how to use a semi-automatic defibrillator.⁹ In addition, while children under 12 years may not be able to deliver quality

chest compressions (possibly on account of anthropometric factors),¹⁰ they are capable of delivering vigorous compressions and learn theoretical concepts.

The *Adestrando Nenos e Xóvenes a Salvar Vidas* project (Teaching children and youth to save lives, ANXOS) addresses the main practical uncertainties that have emerged in relation to school-based BLS training of children and youth. In this context, the aim of our study was to assess the acquisition of BLS theoretical knowledge and skills in students aged less than 13 years through a brief training delivered by their physical education instructors.

Material and methods

Sample

We conducted a quasiexperimental study in which we invited to participate a convenience sample of 658 students in 3 charter schools in Galicia. The study was approved by the Ethics Committee of the School of Education and Sport Sciences of the Universidad de Vigo.

The inclusion criteria were age 8 to 12 years, absence of physical or mental illness and written informed consent of parents and verbal assent of the minor. We excluded participants for who the anthropometric data were not documented or with incomplete data.

Study design

The ANXOS project started by setting up meetings to inform school administrators and parents, explaining the objectives and design of the study, and requesting the necessary authorizations and consent.

The next step was to train physical education instructors,¹¹ who were to be responsible for training students. In a second phase (corresponding to the study presented here), students filled out an ad hoc questionnaire to assess their knowledge a week before undergoing a training structured in 2 sessions that took place in the times allotted to the 2 weekly physical education classes, each of which had a duration of 50 minutes. The first session was devoted to theoretical contents taught with materials adapted to the age of the students; the second session was devoted to practical training with adult-size manikins that provided instant visual feedback on the quality of chest compressions. There was 1 manikin per 5 students and 1 instructor per 25 students. Each student practiced delivering chest compressions on the manikin for a minimum of 6 minutes.

The practical session was organised around 3 stations: 1) the BLS sequence; 2) practice of BLS skills on an adult torso CPR training manikin (without feedback), and 3) practice of BLS skills on the manikin with real-time audio guidance and visual feedback.

After the training sessions, the theoretical knowledge and practical skills of the students were assessed again. Theoretical knowledge was assessed with the same questionnaire used for the baseline assessment at the beginning of the study. We evaluated practical skills through the simulation of a case of OHCA, assessing the ability to follow the

steps of the BLS sequence and the quality of chest compressions delivered over 2 minutes.

Variables

We collected data on the following variables: 1) sociodemographic and anthropometric characteristics (school year, age, weight, height, body mass index); 2) theoretical knowledge, assessed with a questionnaire; 3) BLS sequence: securing the scene, checking for a response, opening the airway, look-listen-feel assessment of breathing, alerting medical emergency services and initiation of chest compressions; 4) CPR quality: time to initiation of compressions (in seconds), chest compression fraction, percentage of compressions with hands in the right position, mean depth of compressions (in mm), percentage of compressions that reached the recommended depth (50–60 mm), percentage of compressions with correct decompression, mean rate of compressions (compressions/minute) and percentage of compressions delivered at the correct rate (100–120/minute).

Assessment tools

Knowledge questionnaire

We designed an ad hoc questionnaire to assess knowledge through 5 items with three answer choices of which only one was correct. The possible score ranged from 0 to 5 points (Appendix B, supplemental file 1).

Basic life support sequence

We assessed the BLS sequence by means of an ad hoc evaluation form that included every step in the sequence and documented the time elapsed from the start of the simulation to initiation of chest compressions (Appendix B, supplemental file 2).

Quality of CPR

To measure the quality of the compression, we used the Laerdal Resusci Anne manikin with SkillReporter (version 12.0.0.2, wireless for PC), setting the parameters according to the 2015 ERC guidelines: chest compression depth of 50–60 mm; compression rate of 100–120 compressions/minute. As a quality criterion, we used a widely accepted arbitrary threshold of 70%.¹²

Statistical analysis

In the analysis of quantitative data, we assessed the normality of the distribution by means of the Kolmogorov-Smirnov test. We summarised quantitative variables using central tendency and dispersion measures (mean/median and standard deviation [SD]/interquartile range [IQR]) and qualitative variables as absolute and relative frequency distributions. We used the Pearson correlation coefficient to analyse the linear association between anthropometric variables and CPR quality variables. The association between categorical variables was assessed by means of the chi square test. We compared means with the Student *t* test or Mann-Whitney *U* test as applicable, and multiple means by




Figure 1 Percentage of participants that performed each step of the basic life support chain sequence post training. Data disaggregated by school year.

analysis of variance (ANOVA) with the Bonferroni correction (homogeneity of variance met) or the Games-Howell correction (homogeneity of variance not met). The data were processed and analysed with the statistical software SPSS version 24.0. We defined statistical significance as a *p*-value less than 0.05.

Results

The initial sample consisted of 658 students, of who we excluded 42 (6.4%) due to incomplete data and 49 (7.4%) due to lack of signed informed consent at the start of the study, so the final sample included 567 students (48.0% female). The distribution by school year was: 32.1% in year 3 of primary education (PE), 31.6% in year 5 of PE, and 36.3% in year 1 of compulsory secondary education (known in Spain as ESO).

Theoretical knowledge

The percentage of students that answered at least 4 questions (out of 5) increased from the pre- to the post-training test in every year: year 3 of PE (2.3% vs 25.6%; $P < .001$), year 5 of PE (1.1% vs 46.2%; $P < .001$) and year 1 of the ESO (23.3% vs 77.0%; $P < .001$).

Basic life support sequence

Fig. 1 and supplemental file 3 in Appendix B present the data concerning the performance of the BLS sequence after the training. At least 50% of the students were able to perform all the steps in the BLS sequence in each school year. The steps the students performed less frequently ($< 70\%$) were "securing the scene" and "checking for a response" (students in year 3 of PE and year 1 of ESO). More than 80% of students performed the "look-listen-feel assessment of breathing", "alerting medical emergency services" and "initiation of chest compressions" steps.

When we analysed the percentage of students that performed the full sequence by school year, we found that a higher proportion of students in year 5 of PE could perform the full sequence compared to year 3 of PE or year 1 of ESO (year 3 of PE: 16.5%; year 5 of PE: 54.4%; year 1 of ESO: 28.5%; $P = .030$). In the subset of students that performed the full sequence, we did not find significant differences based on school year in the seconds elapsed from the start of the simulation to initiation of chest compressions (year 3 of PE: median, 52.2 [IQR, 43.3-58.7]; year 5 of PE: median, 47 [IQR, 42.5-55.3]; year 1 of ESO: median, 40.7 [IQR 40.3-56.4]; $P = .577$).




Figure 2 Overall quality of resuscitation and each compression-related variable. Data disaggregated by school year.

Quality of CPR

Fig. 2 presents the results on the parameters used to assess the overall quality of CPR by school year, with additional results presented in supplemental file 4 in Appendix B.

The chest compression fraction, mean depth of compressions and percentage of compressions with correct depth increased with the school year ($P < .001$). We also found significant linear associations between with CPR quality parameters and anthropometric variables (Table 1).

Fig. 3 presents the results regarding the depth of compressions (mm) and mean rate of compressions (per minute). It shows a target area, marked in red, that includes the recommended values,¹³ and a blue area defined by arbitrary deviation thresholds of ± 10 compressions per minute and ± 5 mm of depth. When it came to depth, the target of 50 mm was only achieved by 3.4% of participants in year 1 of ESO ($P = .002$) compared to none in years 3 and 5 of PE. As for the compression rate, 31.9% of students in year 3 of PE, 33.0% in year 5 of PE and 45.1% in year 1 of ESO maintained an adequate rate of compressions ($P = .003$).

Only 2 students (0.98%) in year 1 of ESO delivered CPR at a depth of 50–60 mm and rate of 100–120 compressions/minute. However, 19 students of year 1 of ESO (9.27%), 4 (2.23%) in year 5 of PE (although none in year 3 of PE) managed to position themselves in the wider arbitrarily defined blue area.

Discussion

Our study shows that following a brief and simple training, delivered during school hours by the physical education instructors of the schools, enabled students in years 3 and 5 of PE and year 1 of ESO (ages 8–12 years) to activate the chain of survival and initiate basic life support, although the quality of CPR was insufficient.

Similar to previous studies, we found that students in the sample were able to assimilate knowledge on first aid and BLS.^{14–16} In fact, the results of a small study in Italy showed that students could even have greater capacity for knowledge acquisition and retention for this type of content compared to adults,¹⁷ which would further justify teaching BLS to schoolchildren without waiting for them to reach adolescence or adulthood.

When we assessed the performance of the BLS sequence, more than half of the children performed each of the steps, a proportion that exceeded 70% in the oldest group. Although most were not able to perform the full sequence with every step in the right order, it is fair to say that they exhibited good knowledge acquisition, as reported in previous studies.^{14,18} It must be taken into account that carrying out each step consecutively in adherence with current guidelines is no simple task even for adults.¹⁹


Another factor at play in the delivery of basic CPR was the physical characteristics of the students, in an age range that is strongly associated with anthropometric growth. Achieving the recommended depth of compression has proven a nearly impossible goal for children under 13 years.^{10,20} This is not the case of other compression quality parameters that are not as dependent on height and weight, such as the chest compression fraction, correct hand positioning or full chest recoil (which happens to be easier for weaker and younger children), for which we observed proportions greater than 85%, as did previous studies.^{10,14,18}

The percentage of compressions followed by adequate decompression decreased with increasing student age. One of the possible explanations is that younger children allow adequate thoracic decompression to occur due to anthropometric factors, as it is more challenging for them to stop the recoil of the manikin's chest. The association between the depth of compression, chest recoil and compression rate has already been demonstrated in the context of out-of-

Table 1 Linear relationship between cardiopulmonary resuscitation variables and anthropometric variables.

	Weight	Height	BMI
	<i>r</i> <i>P</i>	<i>r</i> <i>P</i>	<i>r</i> <i>P</i>
Overall quality of resuscitation	0.572 < .001	0.518 < .001	0.405 < .001
Mean depth of compressions	0.636 < .001	0.618 < .001	0.450 < .001
Percentage of compressions with adequate recoil	−0.237 < .001	−0.167 < .001	−0.228 < .001
Percentage of compressions with adequate depth	0.470 < .001	0.398 < .001	0.334 < .001
Percentage of compressions at adequate rate	0.111 0.008	0.129 0.002	– –

BMI, body mass index.

**Figure 3** Target chest compression rate and depth grid and scatterplot of depth and rate achieved by all participants in each school year. Participants that met the recommended targets are in the red area, those that came close to the targets are in the blue area.

hospital CPR.²¹ When it comes to the compression rate, the factors at play are not only the anthropometric characteristics of rescuers or their ability to assimilate knowledge, but also the level of psychomotor development, which determines the ability to move at a given pace, in this case, at a rate of 100-120 compressions per minute. This explains why, despite the fact that the mean rate of compressions was in the recommended target range in every group, a higher proportion of older children achieved an adequate chest compression fraction. These differences between groups in favour of older students have already been observed in previous studies.^{10,14,22}

The Kids Save Lives statement, issued by the ERC and endorsed by the World Health Organization, has the objective of increasing survival after OHCA and promotes 10 principles, the first of which is that “kids can save lives”.¹

In consequence, the school is considered the ideal setting for training with the aim of having a significant impact on the response to OHCA. In several European countries, BLS training in schools is mandatory and integrated in the academic curriculum²³; in Spain, Royal Decrees 126/2014²⁴ and 1105/2014,²⁵ which establish the primary and secondary education curricula, respectively, include training on first aid, but do not specify the contents to be taught, so that teaching of BLS and CPR is not mandatory as it is in some neighbouring countries.

The inclusion of contents related to first aid in general or BLS in particular requires adequate knowledge of this material by schoolteachers. In a study conducted in Galicia, 70.6% of surveyed teachers reported having been trained on first aid, but the vast majority answered questions about BLS incorrectly.²⁶ Recently, the ERC has issued a position

statement on the subject,²⁷ proposing that once they have been trained, teachers can be as competent as health care professionals in teaching BLS contents. In addition, training by schoolteachers would substantially cut costs compared to having to send health care professionals to each school. Therefore, inclusion of first aid-related subjects in the educational curricula of university courses for the training of early childhood educators and primary education teachers should be considered.^{26,28}

There are limitations to our study. The conditions of the simulation hinder the extrapolation of results to potential real-world victims. On one hand, the psychological variables that affect a real-life situation cannot be introduced in this type of simulated scenarios, and on the other, simulations are conducted on manikins, which, while being a valid and reliable instrument, do not exactly reproduce the conditions of real-world CPR, in which compression parameters are also influenced by the physical characteristics of the victim.

We used only one training method, without a control group, and the assessment was conducted immediately after completion of the training, so that it does not reflect the retention of acquired knowledge and skills in the medium and long terms.

Conclusion

A 2-hour theoretical/practical workshop during school hours and taught by physical education instructors contributes to students learning how to identify an emergency, initiate the chain of survival and initiate basic CPR with chest compressions. The quality of compressions in terms of depth and rhythm proved insufficient. Training on BLS skills should be delivered in every school in Spain, as recommended by the World Health Organization and the ERC.

Conflicts of interest

The authors have no conflicts of interest to declare.

Appendix A. Supplementary data

Supplementary material related to this article can be found, in the online version, at doi:<https://doi.org/10.1016/j.anpede.2020.09.013>.

References

- Böttiger BW, Bossaert LL, Castron M, Cimpoesu D, Georgiou M, Greif R, et al. Kids Save Lives - ERC position statement on school children education in CPR. "Hands that help - Training children is training for life. *Resuscitation*. 2016;105:A1–3.
- Sasson C, Rogers MAM, Dahl J, Kellermann AL. Predictors of survival from out-of-hospital cardiac arrest: A systematic review and meta-analysis. *Circ Cardiovasc Qual Outcomes*. 2010;3:63–81.
- Greif R, Lockey AS, Conaghan P, Lippert A, de Vries W, Mon-sieus KG, et al. European Resuscitation Council Guidelines for Resuscitation 2015: Section 10. Education and implementation of resuscitation. *Resuscitation*. 2015;95:288–301.
- Cerdà M, Chanovas Borrás M, Escalada Roig X, Espuny Vidal C. Plan piloto de formación en soporte vital y RCP en las escuelas. *Med Intensiva*. 2012;36:158–9.
- López Unanua MC, Freire Tellado M, Rasines Sisneaga R, Iglesias González A. RCP na aula: programa de enseñanza de la reanimación cardiopulmonar básica para estudiantes de secundaria. *Emergencias*. 2012;24:74–80.
- Miró Ò, Díaz N, Sánchez M. Aprender reanimación cardiopulmonar desde la escuela. *Emergencias*. 2012;24:423–5.
- Plant N, Taylor K. How best to teach CPR to schoolchildren: A systematic review. *Resuscitation*. 2013;84:415–21.
- Ammirati C, Gagnayre R, Amsellem C, Nemitz B, Gignon M. Are schoolteachers able to teach first aid to children younger than 6 years? A comparative study. *BMJ Open*. 2014;4:e005848.
- Jorge-Soto C, Abelairas-Gómez C, Barcala-Furelos R, Gregorio-García C, Prieto-Saborit JA, Rodríguez-Núñez A. Learning to use semiautomatic external defibrillators through audiovisual materials for schoolchildren. *Emergencias*. 2016;28:103–8.
- Abelairas-Gómez C, Rodríguez-Núñez A, Casillas-Cabana M, Romo-Pérez V, Barcala-Furelos R. Schoolchildren as life savers: At what age do they become strong enough? *Resuscitation*. 2014;85:814–9.
- Pichel López M, Martínez-Isasi S, Barcala-Furelos R, Fernández-Méndez F, Vázquez Santamarina D, Sánchez-Santos L, et al. Un primer paso en la enseñanza del soporte vital básico en las escuelas: la formación de los profesores. *An Pediatr*. 2018;89:265–71.
- Perkins GD, Colquhoun M, Simons R. ABC of resuscitation. 5th ed. London: BMJ Books; 2004. p. 97–101.
- Perkins GD, Handley AJ, Koster RW, Castrén M, Smyth MA, Olasveengen T, et al. European Resuscitation Council Guidelines for Resuscitation 2015: Section 2. Adult basic life support and automated external defibrillation. *Resuscitation*. 2015;95:81–99.
- Banfai B, Pek E, Pandur A, Csonka H, Betlehem J. 'The year of first aid': Effectiveness of a 3-day first aid programme for 7–14-year-old primary school children. *Emerg Med J*. 2017;34:526–32.
- Vetter VL, Haley DM, Dugan NP, Iyer VR, Shults J. Innovative cardiopulmonary resuscitation and automated external defibrillator programs in schools: Results from the Student Program for Olympic Resuscitation Training in Schools (SPORTS) study. *Resuscitation*. 2016;104:46–52.
- Cerezo Espinosa C, Nieto Caballero S, Juguera Rodríguez L, Castejón-Mochón JF, Segura Melgarejo F, Sánchez Martínez CM, et al. Ensayo clínico aleatorizado controlado que compara la formación presencial frente a la no presencial en el aprendizaje teórico de la reanimación cardiopulmonar entre los estudiantes de secundaria. *Emergencias*. 2018;30:28–34.
- Baldi E, Bertaia D. School children learn BLS better and in less time than adults. *Resuscitation*. 2014;85:536–7.
- Li H, Shen X, Xu X, Wang Y, Chu L, Zhao J, et al. Bystander cardiopulmonary resuscitation training in primary and secondary school children in China and the impact of neighborhood socioeconomic status. *Medicine (Baltimore)*. 2018;97:e12673.
- González-Salvado V, Abelairas-Gómez C, Peña-Gil C, Neiro-Rey C, Barcala-Furelos R, González-Juanatey JR, et al. Basic life support training into cardiac rehabilitation programs: A chance to give back. A community intervention controlled manikin study. *Resuscitation*. 2018;127:14–20.
- He D, Huang K, Yang YI, Jiang W, Yang N. What is the optimal age for students to receive cardiopulmonary resuscitation training? *Prehospital Med Med*. 2018;33:394–8.
- González-Otero DM, Russell JK, Ruiz JM, Ruiz de Gauna S, Gutiérrez JJ, Leturiondo LA, et al. Association of chest compression and recoil velocities with depth and rate in manual cardiopulmonary resuscitation. *Resuscitation*. 2019;142:119–26.

22. Jones I, Whitfield R, Colquhoun M, Chamberlain D, Vetter N, Newcombe R. At what age can schoolchildren provide effective chest compressions? An observational study from the Heartstart UK schools training programme. *BMJ*. 2007;334:1201.
23. Semeraro F, Wingen S, Schroeder DC, Ecker H, Scapigliati A, Ristagno G, et al. KIDS SAVE LIVES implementation in Europe: A survey through the ERC Research NET. *Resuscitation*. 2016;107:e7–9.
24. Real Decreto 126/2014, de 28 de febrero, por el que se establece el currículo básico de la Educación Primaria. BOE núm. 52, de 1 de marzo de 2014. p. 19349-19420 [accessed 16 Jun 2020]. Available from: <https://www.boe.es/boe/dias/2014/03/01/pdfs/BOE-A-2014-2222.pdf>.
25. Real Decreto 1105/2014, de 26 de diciembre, por el que se establece el currículo básico de la Educación Secundaria Obligatoria y del Bachillerato [Internet]. BOE núm. 3, de 3 Ene de 2015. p. 169-546 [accessed 16 Jun 2020]. Available from: <https://www.boe.es/boe/dias/2015/01/03/pdfs/BOE-A-2015-37.pdf>.
26. Abelairas-Gómez C, Carballo-Fazanes A, Martínez-Isasi S, López-García S, Rico-Díaz J, Rodríguez-Núñez A. Knowledge and attitudes on first aid and basic life support of pre- and elementary school teachers and parents. *An Pediatr (Barc)*. 2020;92:268–76.
27. Böttiger BW, Lockey A, Georgiou M, Greif R, Monsieurs KG, Mpotos N, et al. KIDS SAVE LIVES: ERC Position statement on schoolteachers' education and qualification in resuscitation. *Resuscitation*. 2020;151:87–90.
28. Abelairas-Gómez C, López-García S, Martínez-Isasi S, Carballo-Fazanes A, Rodríguez-Núñez A. Basic life support knowledge of the future of the Infant and Primary School teacher. An unresolved problem in university study plans? *An Pediatr (Barc)*. 2019;91:344–5.