



## Scientific Letter

### Is the Effect of Occupation on Lung Cancer Modified by Tobacco Use and Residential Radon Exposure? Results of a Pooling Study



To the Director,

Lung cancer is the leading cause of cancer mortality worldwide, accounting for one in every five cancer-related deaths and responsible for approximately 18.7% of all cancer deaths in both sexes [1]. It is common to diagnose lung cancer at advanced stages, impacting its prognosis and survival [2]. The main risk factor associated with this type of cancer is tobacco smoking, however, occupational exposure to carcinogens and radon exposure are also well-known contributors to lung cancer burden. Of the 120 agents classified by the International Agency for Research on Cancer (IARC) as group 1 human carcinogens, 47 are of occupational origin [3]. Such exposures occur in a wide range of occupations, including mining, construction, glass and ceramic fiber work, and chemical manufacturing. Despite this, the evidence regarding occupational risk among women remains limited, since most studies have focused on men and subsequently extrapolated results to women [4–6]. Sex-specific results are therefore necessary.

Radon gas is recognized by the World Health Organization (WHO) as the second leading cause of lung cancer overall, and the leading cause among never-smokers [7]. Radon is a naturally occurring radioactive gas that can accumulate indoor, exposing the respiratory tract to ionizing radiation [8]. The WHO has recommended reference levels of 100 Bq/m<sup>3</sup> [7]. While the interaction between radon and tobacco smoking has been well documented [9–12], evidence on the combined effect of radon, tobacco, and occupational exposures remains scarce [13].

The aim of this study was, first, to determine whether tobacco smoking modifies the effect of occupation on lung cancer risk, stratified by sex; and second, to analyze whether residential radon exposure modifies the effect of occupation on lung cancer risk, again stratified by sex.

We conducted a pooling study of five multicenter case-control studies carried out between 1992 and 2020 in hospitals located in Galicia, Castilla y León, Madrid, Asturias (Spain), and Oporto (Portugal). All original studies used comparable methodologies, allowing harmonization of variables and joint analysis in a single pooled database. Cases were histologically confirmed lung cancer patients, while hospital-based controls were frequency-matched by age and sex. Controls were patients admitted for conditions not related to lung cancer, such as orthopedic surgery, cataracts, lipomas, or inguinal hernias; as well as from uncomplicated microsurgical procedures, including carpal tunnel release or disk hernias. Only the controls from the first study [14] were selected from the 1991 Census, using stratified random sampling based on the sex

distribution among cases and the population weighting of each sub-area.

Trained staff administered structured questionnaires to collect sociodemographic data, tobacco history, occupational history (up to five previous occupations and duration of each occupation), and residential information. Radon exposure was measured in dwellings using alpha-track devices over a minimum of three months, processed at the Galician Radon Laboratory. Of note, 788 participants lacked radon measurements, mainly because the detector was not returned by the participant. To minimize this issue, two follow-up phone calls were made to encourage return of the device. Occupations were coded according to ISCO-68, grouped at the major group level. Additionally, we created a category of “Housewives”.

We first performed logistic regressions stratified by sex, using “Administrative and managerial workers” and “Clerical workers” as reference groups, adjusting for age, smoking, and radon exposure. Odds ratios (ORs) and 95% confidence intervals (95% CIs) were estimated. To explore modification of effect, we fitted models with combined categories of occupation and smoking, and of occupation and radon, globally and by sex. Statistical significance was set at  $p < 0.05$ , and analyses were performed in Stata v.17.

A total of 4203 participants (1987 cases and 2216 controls) were included, with 1288 women and 2915 men. Median age was higher among cases than controls in both sexes. The percentage of never smokers was higher among women (69.8% of female cases, 82.6% of female controls) compared with men (8.0% of male cases, 35.9% of male controls). For ever-smokers, cumulative tobacco consumption was similar in men and women, with cases being heavier smokers compared to controls. Residential radon exposure above 300 Bq/m<sup>3</sup> was slightly more frequent in cases than controls for both sexes. Cases and controls resided mainly in towns and countryside. Adenocarcinoma was the most frequent histologic type in women (63.7%), whereas squamous cell carcinoma predominated in men (29.2%).

In adjusted logistic regression models, several occupational categories were associated with increased lung cancer risk, with differences by sex. Among women, “Sales workers” (OR 5.19, 95%CI: 1.79–15.08), “Craft and related trades workers” (OR 1.96, 95%CI: 1.04–3.69), “Elementary occupations” (OR 5.12, 95%CI: 1.26–20.86), and “Housewives” (OR 2.61, 95%CI: 1.47–4.62) showed significant ORs. Among men, significant associations were observed for “Professional, technical and related workers” (OR 1.79, 95%CI: 1.10–2.90), “Service workers” (OR 1.65, 95%CI: 1.06–2.55), “Craft and related trades workers” (OR 1.70, 95%CI: 1.05–2.75), “Plant and machine operators” (OR 1.78, 95%CI: 1.19–2.65), and “Elementary occupations” (OR 1.69, 95%CI: 1.18–2.41).

When analyzing the joint effect of tobacco smoking and occupation, we observed a possible modification effect among men.

**Table 1**  
Tobacco use, major occupational groups (ISCO-68) and lung cancer risk, stratified by sex.

ISCO-68	Major groups	Never smokers			Exsmokers			Smokers		
		n	OR	95%CI	n	OR	95%CI	n	OR	95%CI
<i>Women</i>										
2/3	Administrative, managerial, clerical and related workers	42	1	–	25	1.18	0.41–3.41	24	1.32	0.43–4.04
0/1	Professional, technical and related workers	88	0.85	0.38–1.89	30	2.27	0.84–6.14	32	7.75	2.51–23.86
5	Service workers	236	0.81	0.39–1.65	44	1.19	0.47–3.02	64	3.44	1.42–8.36
6	Agricultural, animal husbandry and forestry workers	250	0.86	0.42–1.79	3	1	–	3	1	–
7	Craft and related trades workers	98	1.37	0.62–3.01	11	1.44	0.32–6.39	4	5.45	0.51–58.60
XI	Housewives/homemakers	201	1.59	0.77–3.27	9	1	–	11	16.66	1.84–150.83
XII	Other	19	7.30	1.41–37.83	11	7.12	1.27–39.97	9	12.66	1.39–115.30
<i>Men</i>										
2/3	Administrative, managerial, clerical and related workers	47	1	–	109	5.13	1.63–16.16	71	17.00	5.19–55.66
0/1	Professional, technical and related workers	41	3.80	1.01–14.27	101	9.32	2.95–29.48	32	22.89	6.01–87.22
4	Sales workers	23	3.55	0.84–14.95	41	18.18	4.93–67.04	26	51.20	10.72–244.48
5	Service workers	59	3.14	0.90–11.00	129	7.29	2.35–22.64	93	29.28	8.81–97.38
6	Agricultural, animal husbandry and forestry workers	175	0.87	0.27–2.83	228	6.49	2.15–19.63	140	8.29	2.69–25.52
7	Craft and related trades workers	41	3.72	1.03–13.40	96	6.45	2.02–20.59	59	37.54	10.11–139.35
8	Plant and machine operators	99	2.80	0.86–9.07	186	10.76	3.54–32.66	111	19.25	6.11–60.68
9	Elementary occupations	166	1.26	0.39–4.07	472	11.97	4.07–35.22	309	19.95	6.68–59.59
XII	Other	10	7.48	1.18–47.34	30	18.48	4.30–79.45	13	47.87	8.30–276.01

Results shown for groups having more than 30 participants in at least one stratum.

**Table 2**  
Residential radon, major occupational groups (ISCO-68) and lung cancer risk, stratified by sex.

ISCO-68	Major groups	<100 Bq/m <sup>3</sup>			100–300 Bq/m <sup>3</sup>			>300 Bq/m <sup>3</sup>		
		n	OR	95%CI	n	OR	95%CI	n	OR	95%CI
<i>Women</i>										
2/3	Administrative, managerial, clerical and related workers	28	1	–	39	1.22	0.43–3.43	17	0.26	0.06–1.17
0/1	Professional, technical and related workers	44	2.34	0.85–6.45	70	1.24	0.48–3.16	25	0.75	0.22–2.53
5	Service workers	23	0.98	0.39–2.46	149	0.99	0.41–2.39	52	1.30	0.49–3.47
6	Agricultural, animal husbandry and forestry workers	55	0.93	0.34–2.52	109	1.04	0.42–2.62	41	2.24	0.78–6.39
7	Craft and related trades workers	18	0.96	0.27–3.40	62	1.51	0.58–3.93	18	5.77	1.42–23.37
XI	Housewives/homemakers	81	1.87	0.74–4.74	84	2.29	0.90–5.80	32	3.58	1.14–11.21
<i>Men</i>										
2/3	Administrative, managerial, clerical and related workers	73	1	–	90	0.68	0.34–1.37	29	1.19	0.45–3.13
0/1	Professional, technical and related workers	52	1.85	0.83–4.10	67	1.59	0.75–3.37	27	0.87	0.31–2.47
4	Sales workers	37	1.09	0.45–2.65	27	7.76	2.49–24.24	7	5.58	0.76–41.12
5	Service workers	99	1.47	0.74–2.90	75	1.49	0.72–3.09	39	0.84	0.35–2.01
6	Agricultural, animal husbandry and forestry workers	273	0.43	0.23–0.78	115	1.24	0.64–2.43	40	0.99	0.41–2.36
7	Craft and related trades workers	69	0.93	0.44–1.95	65	1.55	0.73–3.31	17	5.80	1.46–23.05
8	Plant and machine operators	138	1.09	0.57–2.08	126	1.93	1.00–3.70	57	1.61	0.74–3.52
9	Elementary occupations	371	1.04	0.59–1.83	270	1.72	0.96–3.10	112	1.77	0.91–3.46

Results shown for groups having more than 30 participants in at least one stratum.

For instance, in men, “Professional, technical and related workers” showed ORs rising from 3.80 (95%CI: 1.01–14.27) among never-smokers to 22.89 (95%CI: 6.01–87.22) among smokers. Similarly, “Craft and related trades workers” presented an OR of 3.72 (95%CI: 1.03–13.40) among never-smokers, which increased to 37.54 (95%CI: 10.11–139.35) among smokers. Among women, this effect was less consistent, with some significant results such as “Service workers” (OR 3.44, 95%CI: 1.42–8.36) and “Housewives” (OR 16.66, 95%CI: 1.84–150.83) among smokers, but overall weaker trends (Table 1).

Regarding radon exposure, effect modification by occupation was suggested for both sexes, however, results at lower exposure levels were not statistically significant. Among women exposed to >300 Bq/m<sup>3</sup>, risks were particularly high for “Craft and related trades workers” (OR 5.77, 95%CI: 1.42–23.37) and “Housewives” (OR 3.58, 95%CI: 1.14–11.21). Among men in the same exposure category, “Craft and related trades workers” showed an OR of 5.80 (95%CI: 1.46–23.05) (Table 2).

Our findings suggest that both tobacco smoking and residential radon exposure may modify the effect of occupational exposures on lung cancer risk, particularly among men. To our knowledge, this is the first study to assess potential modification between occupa-

tion and radon exposure in a general population. Previous research on miners has described submultiplicative interactions between radon and smoking [15,16], whereas our data indicate that residential radon could have an independent and possibly additive effect in the general population.

Sex differences observed in this study may reflect not only biological factors, such as hormonal influences and endocrine interactions with tobacco [17,18], but also differences in occupational distribution between men and women. Men predominate in heavy industrial trades, while women are more frequently represented in service jobs, where exposures may differ. These results align with previous reports, highlighting the importance of considering sex in occupational cancer epidemiology [4,5].

The main limitation of this work lies in the use of ISCO-68 major groups, which pool heterogeneous occupations within broad categories. This may mask risks specific to certain jobs or industries. Additionally, our analysis was based on occupational categories rather than direct assessment of occupational exposures. A more descriptive analysis linking specific exposures with lung cancer risk would provide additional results. Interaction models were not conducted due to sample size limitations, so we opted for stratified analyses as an exploratory approach. Future studies with larger

sample sizes will be needed to evaluate interaction terms with greater statistical power. Finally, stratification by sex, radon, and smoking inevitably reduced sample sizes in some subgroups, limiting statistical power. Strengths of this study include its large sample size, adjustment for residential radon exposure (not usually considered in previous studies) and sex-stratified analyses.

In conclusion, our study indicates that occupational risk of lung cancer may be modified by tobacco smoking and, to a lesser extent, by residential radon exposure. These effects differ by sex, with stronger associations observed among men. Our results highlight the need for preventive strategies such as smoking cessation programs directed to workers exposed to occupational carcinogens, radon mitigation in radon-prone areas and the consideration of occupational history into lung cancer screening.

### Author contribution

Eliana Torres-Cadavid: conceptualization, resources, data curation, writing – original draft. Cristina Candal-Pedreira: conceptualization, methodology, formal analysis, investigation, writing – review & editing. Alberto Ruano-Ravina: Conceptualization, methodology, data curation, supervision, writing – review & editing. Karl Kelsey: writing – review & editing. Lucía Martín-Gisbert: investigation, writing – review & editing.

### Ethics approval

The original studies protocols, funding and informed consent models were approved by the relevant Clinical Research Ethics Committees.

### Declaration

This scientific letter forms part of the work leading to the doctoral thesis of Eliana Torres Cadavid.

### Artificial intelligence involvement

No artificial intelligence tools were used in process of this study. All work was carried out by the authors.

### Funding

No funding was received.

### Conflicts of interest

Dr. Kelsey is a founder and scientific advisor to Cellintec which had no role in this research. The remaining authors declare there are no conflicts of interest that are relevant to the publication of this paper.

### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi: 10.1016/j.arbres.2025.10.009](https://doi.org/10.1016/j.arbres.2025.10.009).

### References

- [1] Bray F, Laversanne M, Sung H, Ferlay J, Siegel RL, Soerjomataram I, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2024;74(3):229–63.
- [2] Allemani C, Matsuda T, Di Carlo V, Harewood R, Matz M, Nikšić M, et al. Global surveillance of trends in cancer survival 2000–14 (CONCORD-3): analysis of individual records for 37,513,025 patients diagnosed with one of 18 cancers from 322 population-based registries in 71 countries. *Lancet Lond Engl* 2018;391(10125):1023–75.
- [3] Loomis D, Guha N, Hall AL, Straif K. Identifying occupational carcinogens: an update from the IARC Monographs. *Occup Environ Med* 2018;75(8):593–603.
- [4] Betansedi CO, Vaca Vasquez P, Counil E. A comprehensive approach of the gender bias in occupational cancer epidemiology: a systematic review of lung cancer studies (2003–2014). *Am J Ind Med* 2018;61(5):372–82.
- [5] Stapelfeld C, Dammann C, Maser E. Sex-specificity in lung cancer risk. *Int J Cancer* 2020;146(9):2376–82.
- [6] Siegfried JM. Sex and gender differences in lung cancer and chronic obstructive lung disease. *Endocrinology* 2022;163(2):bqab254.
- [7] World Health Organization. WHO handbook on indoor radon: a public health perspective. Geneva: World Health Organization; 2009. p. 111.
- [8] Henyoh AMS, Laurent O, Mandin C, Clero E. Radon exposure and potential health effects other than lung cancer: a systematic review and meta-analysis. *Front Public Health* 2024;12:1439355.
- [9] Ruano-Ravina A, Martín-Gisbert L, Kelsey K, Pérez-Ríos M, Candal-Pedreira C, Rey-Brandariz J, et al. An overview on the relationship between residential radon and lung cancer: what we know and future research. *Clin Transl Oncol* 2023;25(12):3357–68.
- [10] Cheng ES, Egger S, Hughes S, Weber M, Steinberg J, Rahman B, et al. Systematic review and meta-analysis of residential radon and lung cancer in never-smokers. *Eur Respir Rev* 2021;30(159):200230.
- [11] Park EJ, Lee H, Kim HC, Sheen SS, Koh SB, Park KS, et al. Residential radon exposure and cigarette smoking in association with lung cancer: a matched case-control study in Korea. *Int J Environ Res Public Health* 2020;17(8):2946.
- [12] Tomasek L. Lung cancer risk from occupational and environmental radon and role of smoking in two Czech nested case-control studies. *Int J Environ Res Public Health* 2013;10(3):963–79.
- [13] Grandi C, S anjust F. 124 occupational exposure to radon: an underestimated risk in view of the combined exposure to other occupational and environmental lung carcinogens. *Occup Environ Med* 2018;75 Suppl. 2. A426.2.
- [14] Barros-Dios JM, Barreiro MA, Ruano-Ravina A, Figueiras A. Exposure to residential radon and lung cancer in Spain: a population-based case-control study. *Am J Epidemiol* 2002;156(6):548–55.
- [15] Schubauer-Berigan MK, Bertke SJ, Kelly-Reif K, Daniels RD. Updated cancer mortality among uranium miners on the Colorado Plateau: interactions of radon exposure with smoking and temporal factors. *Occup Environ Med* 2025;82(5):230–7.
- [16] Leuraud K, Schnelzer M, Tomasek L, Hunter N, Timarche M, Grosche B, et al. Radon, smoking and lung cancer risk: results of a joint analysis of three European case-control studies among uranium miners. *Radiat Res* 2011;176(3):375–87.
- [17] Townsend EA, Miller VM, Prakash YS. Sex differences and sex steroids in lung health and disease. *Endocr Rev* 2012;33(1):1–47.
- [18] Rodríguez-Lara V, Avila-Costa MR. An overview of lung cancer in women and the impact of estrogen in lung carcinogenesis and lung cancer treatment. *Front Med (Lausanne)* 2021;8:600121.

Eliana Torres-Cadavid<sup>a</sup>, Cristina Candal-Pedreira<sup>a,b,c,\*</sup>,  
Alberto Ruano-Ravina<sup>a,b,c</sup>, Karl Kelsey<sup>d</sup>,  
Lucía Martín-Gisbert<sup>a,e</sup>

<sup>a</sup> Department of Preventive Medicine and Public Health, University of Santiago de Compostela, Santiago de Compostela, Spain

<sup>b</sup> Consortium for Biomedical Research in Epidemiology and Public Health (CIBER en Epidemiología y Salud Pública, CIBERESP), Spain

<sup>c</sup> Health Research Institute of Santiago de Compostela (Instituto de Investigación Sanitaria de Santiago de Compostela – IDIS), Santiago de Compostela, Spain

<sup>d</sup> Department of Epidemiology, Brown School of Public Health, Brown University, Providence, United States

<sup>e</sup> Cross-disciplinary Research in Environmental Technologies (CRETUS), University of Santiago de Compostela, Santiago de Compostela, Spain

Corresponding author.

E-mail address: [c.candal.pedreira@usc.es](mailto:c.candal.pedreira@usc.es) (C. Candal-Pedreira).