

Research Article

On a New Class of Antiperiodic Fractional Boundary Value Problems

Bashir Ahmad,¹ Juan J. Nieto,^{1,2} Ahmed Alsaedi,¹ and Nadia Mohamad¹

¹ Department of Mathematics, Faculty of Science, King Abdulaziz University, P.O. Box 80203, Jeddah 21589, Saudi Arabia

² Departamento de Análisis Matemático, Facultad de Matemáticas, Universidad de Santiago de Compostela, 15782 Santiago de Compostela, Spain

Correspondence should be addressed to Juan J. Nieto; juanjose.nieto.roig@usc.es

Received 9 October 2012; Accepted 6 February 2013

Academic Editor: Ağacık Zafer

Copyright © 2013 Bashir Ahmad et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

This paper investigates a new class of antiperiodic boundary value problems of higher order fractional differential equations. Some existence and uniqueness results are obtained by applying some standard fixed point principles. Some examples are given to illustrate the results.

1. Introduction

Boundary value problems of fractional differential equations involving a variety of boundary conditions have recently been investigated by several researchers. It has been mainly due to the occurrence of fractional differential equations in a number of disciplines such as physics, chemistry, biology, economics, control theory, signal and image processing, biophysics, blood flow phenomena, aerodynamics, and fitting of experimental data. For details and examples, see [1–5]. The recent development of the subject can be found, for example, in papers [6–16].

The mathematical modeling of a variety of physical processes gives rise to a class of antiperiodic boundary value problems. This class of problems has recently received considerable attention; for instance, see [17–24] and the references therein. In [22], the authors studied a Caputo-type antiperiodic fractional boundary value problem of the form

$$\begin{aligned} {}^c D^q x(t) &= f(t, x(t)), \quad t \in [0, T], \quad T > 0, \quad 1 < q \leq 2, \\ x(0) &= -x(T), \end{aligned} \quad (1)$$

$${}^c D^p x(0) = -{}^c D^p x(T), \quad 0 < p < 1.$$

In this paper, we investigate a new class of antiperiodic fractional boundary value problems given by

$$\begin{aligned} {}^c D^q x(t) &= f(t, x(t)), \quad t \in [0, T], \quad T > 0, \quad 2 < q \leq 3, \\ x(0) &= -x(T), \end{aligned}$$

$$\begin{aligned} {}^c D^p x(0) &= -{}^c D^p x(T), \\ {}^c D^{p+1} x(0) &= -{}^c D^{p+1} x(T), \quad 0 < p < 1, \end{aligned} \quad (2)$$

where ${}^c D^q$ denotes the Caputo fractional derivative of order q and f is a given continuous function. Some new existence and uniqueness results are obtained for problem (2) by using standard fixed point theorems.

2. Preliminaries

Let us recall some basic definitions [1–3].

Definition 1. The Riemann-Liouville fractional integral of order q for a continuous function $g : [0, +\infty) \rightarrow \mathbb{R}$ is defined as

$$I^q g(t) = \frac{1}{\Gamma(q)} \int_0^t \frac{g(s)}{(t-s)^{1-q}} ds, \quad q > 0, \quad (3)$$

provided the integral exists.

Definition 2. For $(n-1)$ times absolutely continuous function $g : [0, +\infty) \rightarrow \mathbb{R}$, the Caputo derivative of fractional order q is defined as

$${}^c D^q g(t) = \frac{1}{\Gamma(n-q)} \int_0^t (t-s)^{n-q-1} g^{(n)}(s) ds, \quad (4)$$

$$n-1 < q < n, \quad n = [q] + 1,$$

where $[q]$ denotes the integer part of the real number q .

Notice that the Caputo derivative of a constant is zero.

Lemma 3. For any $y \in C[0, 1]$, the unique solution of the linear fractional boundary value problem

$$\begin{aligned} {}^c D^q x(t) &= y(t), \quad 0 < t < T, \quad 2 < q \leq 3, \\ x(0) &= -x(T), \\ {}^c D^p x(0) &= -{}^c D^p x(T), \\ {}^c D^{p+1} x(0) &= -{}^c D^{p+1} x(T), \quad 0 < p < 1, \end{aligned} \tag{5}$$

is

$$x(t) = \int_0^T G_T(t, s) y(s) ds, \tag{6}$$

where $G_T(t, s)$ is Green's function (depending on q and p) given by

$$G_T(t, s) = \begin{cases} \frac{2(t-s)^{q-1} - (T-s)^{q-1}}{2\Gamma(q)} + \frac{\Gamma(2-p)(T-2t)(T-s)^{q-p-1}}{2T^{1-p}\Gamma(q-p)} + \frac{2T^{1-p}\Gamma(q-p)}{(\Gamma(2-p))^2 T^{p-1}(T-s)^{q-p-2}} + \frac{4\Gamma(q-p-1)\Gamma(3-p)}{(T^2-2t^2)\Gamma(3-p)} - 2T^2 + 4tT, & s \leq t, \\ -\frac{(T-s)^{q-1}}{2\Gamma(q)} + \frac{\Gamma(2-p)(T-2t)(T-s)^{q-p-1}}{2T^{1-p}\Gamma(q-p)} + \frac{(\Gamma(2-p))^2 T^{p-1}(T-s)^{q-p-2}}{4\Gamma(q-p-1)\Gamma(3-p)} + \frac{4\Gamma(q-p-1)\Gamma(3-p)}{(T^2-2t^2)\Gamma(3-p)} - 2T^2 + 4tT, & t < s. \end{cases} \tag{7}$$

Proof. We know that the general solution of equation ${}^c D^q x(t) = y(t)$, $2 < q \leq 3$ can be written as [3]

$$x(t) = \int_0^t \frac{(t-s)^{q-1}}{\Gamma(q)} y(s) ds - b_0 - b_1 t - b_2 t^2, \tag{8}$$

for some constants b_0, b_1 , and $b_2 \in \mathbb{R}$. Using the facts ${}^c D^p b = 0$ (b is a constant),

$$\begin{aligned} {}^c D^p t &= \frac{t^{1-p}}{\Gamma(2-p)}, & {}^c D^p t^2 &= \frac{2t^{2-p}}{\Gamma(3-p)}, \\ {}^c D^{p+1} t^2 &= \frac{2t^{1-p}}{\Gamma(2-p)}, & {}^c D^p I^q y(t) &= I^{q-p} y(t), \end{aligned} \tag{9}$$

we get

$$\begin{aligned} {}^c D^p x(t) &= \int_0^t \frac{(t-s)^{q-p-1}}{\Gamma(q-p)} y(s) ds \\ &\quad - b_1 \frac{t^{1-p}}{\Gamma(2-p)} - \frac{2b_2 t^{2-p}}{\Gamma(3-p)}, \\ {}^c D^{p+1} x(t) &= \int_0^t \frac{(t-s)^{q-p-2}}{\Gamma(q-p-1)} y(s) ds - 2b_2 \frac{t^{1-p}}{\Gamma(2-p)}. \end{aligned} \tag{10}$$

Applying the boundary conditions for the problem (5), we find that

$$\begin{aligned} b_0 &= \frac{1}{2\Gamma(q)} \int_0^T (T-s)^{q-1} y(s) ds \\ &\quad - \frac{T^p \Gamma(2-p)}{2} \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} y(s) ds \\ &\quad + \frac{T^{p+1} \Gamma(2-p)}{2} \left\{ \frac{\Gamma(2-p)}{\Gamma(3-p)} - \frac{1}{2} \right\} \\ &\quad \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} y(s) ds, \\ b_1 &= \Gamma(2-p) T^{p-1} \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} y(s) ds \\ &\quad - \frac{(\Gamma(2-p))^2 T^p}{\Gamma(3-p)} \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} y(s) ds, \\ b_2 &= \frac{\Gamma(2-p)}{2T^{1-p}} \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} y(s) ds. \end{aligned} \tag{11}$$

Substituting the values of b_0, b_1 , and b_2 in (8), we get the solution (6). This completes the proof. \square

Remark 4. For $p = 1$, the solution of the antiperiodic problem

$$\begin{aligned} {}^c D^q x(t) &= y(t), & x(0) &= -x(T), \\ x'(0) &= -x'(T), & x''(0) &= -x''(T), \\ & & 0 < t < T, & 2 < q \leq 3, \end{aligned} \tag{12}$$

is given by [18]

$$x(t) = \int_0^T g(t, s) y(s) ds, \tag{13}$$

where $g(t, s)$ is

$$g(t, s) = \begin{cases} \frac{(t-s)^{q-1} - (1/2)(T-s)^{q-1}}{\Gamma(q)} + \frac{(T-2t)(T-s)^{q-2}}{4\Gamma(q-1)} + \frac{t(T-t)(T-s)^{q-3}}{4\Gamma(q-2)}, & s \leq t, \\ -\frac{(T-s)^{q-1}}{2\Gamma(q)} + \frac{(T-2t)(T-s)^{q-2}}{4\Gamma(q-1)} + \frac{t(T-t)(T-s)^{q-3}}{4\Gamma(q-2)}, & t < s. \end{cases} \quad (14)$$

If we let $p \rightarrow 1^-$ in (7), we obtain

$$G_T(t, s)|_{p \rightarrow 1^-} = \begin{cases} \frac{(t-s)^{q-1} - (1/2)(T-s)^{q-1}}{\Gamma(q)} + \frac{(T-2t)(T-s)^{q-2}}{2\Gamma(q-1)} + \frac{(-2t^2 - T^2 + 4tT)(T-s)^{q-3}}{4\Gamma(q-2)}, & s \leq t, \\ -\frac{(T-s)^{q-1}}{2\Gamma(q)} + \frac{(T-2t)(T-s)^{q-2}}{2\Gamma(q-1)} + \frac{(-2t^2 - T^2 + 4tT)(T-s)^{q-3}}{4\Gamma(q-2)}, & t < s. \end{cases} \quad (15)$$

We note that the solutions given by (14) and (15) are different. As a matter of fact, (15) contains an additional term: $(-t^2 - T^2 + 3tT)(T-s)^{q-3}/4\Gamma(q-2)$. Therefore the fractional boundary conditions introduced in (2) give rise to a new class of problems.

Remark 5. When the phenomenon of antiperiodicity occurs at an intermediate point $\eta \in (0, T)$, the parametric-type antiperiodic fractional boundary value problem takes the form

$$\begin{aligned} {}^c D^q x(t) &= f(t, x(t)), \quad t \in [0, T], \quad 2 < q \leq 3, \\ x(0) &= -x(\eta), \quad {}^c D^p(0) = -{}^c D^p(\eta), \\ {}^c D^{p+1}(0) &= -{}^c D^{p+1}(\eta), \end{aligned} \quad (16)$$

whose solution is

$$x(t) = \int_0^T G_\eta(t, s) f(s, x(s)) ds, \quad (17)$$

where $G_\eta(t, s)$ is given by (7). Notice that $G_\eta(t, s) \rightarrow G_T(t, s)$ when $\eta \rightarrow T^-$.

3. Existence Results

Let $\mathfrak{C} = C([0, T], R)$ denotes a Banach space of all continuous functions defined on $[0, T]$ into R endowed with the usual supremum norm.

In relation to (2), we define an operator $\mathcal{F} : \mathfrak{C} \rightarrow \mathfrak{C}$ as

$$\begin{aligned} (\mathcal{F}x)(t) &= \int_0^t \frac{(t-s)^{q-1}}{\Gamma(q)} f(s, x(s)) ds \\ &\quad - \frac{1}{2} \int_0^T \frac{(T-s)^{q-1}}{\Gamma(q)} f(s, x(s)) ds \\ &\quad + \frac{\Gamma(2-p)(T-2t)}{2T^{1-p}} \\ &\quad \times \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} f(s, x(s)) ds \\ &\quad + \Gamma(2-p)T^{p-1} \\ &\quad \times \left(T^2 - 2t^2 - \frac{2T^2\Gamma(2-p)}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right) \\ &\quad \times (4)^{-1} \\ &\quad \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} f(s, x(s)) ds, \quad t \in [0, T]. \end{aligned} \quad (18)$$

Observe that the problem (2) has a solution if and only if the operator \mathcal{F} has a fixed point.

For the sequel, we need the following known fixed point theorems.

Theorem 6 (see [25]). *Let X be a Banach space. Assume that $T : X \rightarrow X$ is a completely continuous operator and the set $V = \{u \in X \mid u = \mu Tu, 0 < \mu < 1\}$ is bounded. Then T has a fixed point in X .*

Theorem 7 (see [25]). *Let X be a Banach space. Assume that Ω is an open bounded subset of X with $\theta \in \Omega$ and let $T : \overline{\Omega} \rightarrow X$ be a completely continuous operator such that*

$$\|Tu\| \leq \|u\|, \quad \forall u \in \partial\Omega. \quad (19)$$

Then T has a fixed point in $\overline{\Omega}$.

Now we are in a position to present the main results of the paper.

Theorem 8. *Assume that there exists a positive constant L_1 such that $|f(t, x(t))| \leq L_1$ for $t \in [0, T]$, $x \in \mathfrak{C}$. Then the problem (2) has at least one solution.*

Proof. First, we show that the operator \mathcal{F} is completely continuous. Clearly continuity of the operator \mathcal{F} follows from the continuity of f . Let $\Omega \subset \mathfrak{C}$ be bounded. Then, for

all $x \in \Omega$ together with the assumption $|f(t, x(t))| \leq L_1$, we get

$$\begin{aligned}
& |(\mathcal{F}x)(t)| \\
& \leq \int_0^t \frac{(t-s)^{q-1}}{\Gamma(q)} |f(s, x(s))| ds \\
& \quad + \frac{1}{2} \int_0^T \frac{(T-s)^{q-1}}{\Gamma(q)} |f(s, x(s))| ds \\
& \quad + \frac{\Gamma(2-p)|T-2t|}{2T^{1-p}} \\
& \quad \times \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} |f(s, x(s))| ds \\
& \quad + \Gamma(2-p)T^{p-1} \\
& \quad \times \left| T^2 - 2t^2 - \frac{2\Gamma(2-p)T^2}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right| (4)^{-1} \\
& \quad \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} |f(s, x(s))| ds \\
& \leq L_1 \left\{ \int_0^t \frac{(t-s)^{q-1}}{\Gamma(q)} ds + \frac{1}{2} \int_0^T \frac{(T-s)^{q-1}}{\Gamma(q)} ds \right. \\
& \quad \left. + \frac{\Gamma(2-p)}{2T^{1-p}} |T-2t| \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} ds \right. \\
& \quad \left. + \Gamma(2-p)T^{p-1} \right. \\
& \quad \times \left| T^2 - 2t^2 - \frac{2\Gamma(2-p)T^2}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right| (4)^{-1} \\
& \quad \left. \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} ds \right\} \\
& \leq L_1 \left\{ \frac{3T^q}{2\Gamma(q+1)} + \frac{\Gamma(2-p)T^q}{2\Gamma(q-p+1)} + \frac{\Gamma(2-p)T^q}{4\Gamma(q-p)} \right. \\
& \quad \left. \times \left(1 - \frac{2\Gamma(2-p)}{\Gamma(3-p)} + 2 \left(\frac{\Gamma(2-p)}{\Gamma(3-p)} \right)^2 \right) \right\} = M_1, \tag{20}
\end{aligned}$$

which implies that $\|(\mathcal{F}x)(t)\| \leq M_1$.

Furthermore,

$$\begin{aligned}
& |(\mathcal{F}x)'(t)| \\
& \leq \int_0^t \frac{(t-s)^{q-2}}{\Gamma(q-1)} |f(s, x(s))| ds \\
& \quad + \Gamma(2-p)T^{p-1} \\
& \quad \times \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} |f(s, x(s))| ds
\end{aligned}$$

$$\begin{aligned}
& + \frac{\Gamma(2-p)T^{p-1}}{4} \left| \frac{4T\Gamma(2-p)}{\Gamma(3-p)} - 4t \right| \\
& \quad \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} |f(s, x(s))| ds \\
& \leq L_1 T^{q-1} \left\{ \left| \frac{1}{\Gamma(q)} + \frac{\Gamma(2-p)}{\Gamma(q-p+1)} \right. \right. \\
& \quad \left. \left. \times \left\{ 1 + \frac{(q-p)\Gamma(2-p)}{\Gamma(3-p)} \right\} \right| \right\} = M_2. \tag{21}
\end{aligned}$$

Hence, for $t_1, t_2 \in [0, T]$, we have

$$\begin{aligned}
& |(\mathcal{F}x)(t_2) - (\mathcal{F}x)(t_1)| \leq \int_{t_1}^{t_2} |(\mathcal{F}x)'(s)| ds \\
& \leq M_2(t_2 - t_1). \tag{22}
\end{aligned}$$

This implies that \mathcal{F} is equicontinuous on $[0, T]$, by the Arzela-Ascoli theorem, the operator $\mathcal{F} : \mathfrak{C} \rightarrow \mathfrak{C}$ is completely continuous.

Next, we consider the set

$$V = \{x \in \mathfrak{C} \mid x = \mu \mathcal{F}x, 0 < \mu < 1\} \tag{23}$$

and show that the set V is bounded. Let $x \in V$, then $x = \mu \mathcal{F}x, 0 < \mu < 1$. For any $t \in [0, T]$, we have

$$\begin{aligned}
x(t) &= \int_0^t \frac{(t-s)^{q-1}}{\Gamma(q)} f(s, x(s)) ds \\
& \quad - \frac{1}{2} \int_0^T \frac{(T-s)^{q-1}}{\Gamma(q)} f(s, x(s)) ds \\
& \quad + \frac{\Gamma(2-p)(T-2t)}{2T^{1-p}} \\
& \quad \times \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} f(s, x(s)) ds \\
& \quad + \Gamma(2-p)T^{p-1} \\
& \quad \times \left(T^2 - 2t^2 - \frac{2T^2\Gamma(2-p)}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right) (4)^{-1} \\
& \quad \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} f(s, x(s)) ds,
\end{aligned}$$

$$\begin{aligned}
 |x(t)| &= \mu |(\mathcal{F}x)(t)| \\
 &\leq \int_0^t \frac{(t-s)^{q-1}}{\Gamma(q)} |f(s, x(s))| ds \\
 &\quad + \frac{1}{2} \int_0^T \frac{(T-s)^{q-1}}{\Gamma(q)} |f(s, x(s))| ds \\
 &\quad + \frac{\Gamma(2-p)|T-2t|}{2T^{1-p}} \\
 &\quad \times \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} |f(s, x(s))| ds \\
 &\quad + \Gamma(2-p)T^{p-1} \\
 &\quad \times \left| T^2 - 2t^2 - \frac{2\Gamma(2-p)T^2}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right| (4)^{-1} \\
 &\quad \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} |f(s, x(s))| ds \\
 &\leq L_1 \max_{t \in [0, T]} \left\{ \frac{2t^q + T^q}{2\Gamma(q+1)} \right. \\
 &\quad + \frac{\Gamma(2-p)|T-2t|T^{q-1}}{2\Gamma(q-p+1)} \\
 &\quad + \Gamma(2-p)T^{q-2} \\
 &\quad \times \left| T^2 - 2t^2 - \frac{2\Gamma(2-p)T^2}{\Gamma(3-p)} \right. \\
 &\quad \left. \left. + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right| (4)^{-1} \right\} = M_3.
 \end{aligned} \tag{24}$$

Thus, $\|x\| \leq M_3$ for any $t \in [0, T]$. So, the set V is bounded. Thus, by the conclusion of Theorem 6, the operator \mathcal{F} has at least one fixed point, which implies that (2) has at least one solution. \square

Theorem 9. *Let there exists a positive constant r such that $|f(t, x)| \leq \delta|x|$ with $0 < |x| < r$, where δ is a positive constant satisfying*

$$\begin{aligned}
 \max_{t \in [0, T]} \left\{ \frac{2|t|^q + T^q}{2\Gamma(q+1)} + \frac{\Gamma(2-p)T^{q-1}|T-2t|}{2\Gamma(q-p+1)} \right. \\
 + \frac{\Gamma(2-p)T^{q-2}}{4\Gamma(q-p)} \\
 \left. \times \left| T^2 - 2t^2 - \frac{2\Gamma(2-p)T^2}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right| \right\} \delta \leq 1.
 \end{aligned} \tag{25}$$

Then the problem (2) has at least one solution.

Proof. Define $\Omega_1 = \{x \in \mathfrak{C} : \|x\| < r\}$ and take $x \in \mathfrak{C}$ such that $\|x\| = r$; that is, $x \in \partial\Omega$. As before, it can be shown that \mathcal{F} is completely continuous and that

$$\begin{aligned}
 |\mathcal{F}x(t)| \\
 \leq \max_{t \in [0, T]} \left\{ \frac{2|t|^q + T^q}{2\Gamma(q+1)} \right. \\
 + \frac{\Gamma(2-p)T^{q-1}|T-2t|}{2\Gamma(q-p+1)} \\
 + \frac{\Gamma(2-p)T^{q-2}}{4\Gamma(q-p)} \\
 \left. \times \left| T^2 - 2t^2 - \frac{2\Gamma(2-p)T^2}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right| \right\} \\
 \times \delta \|x\| \leq \|x\|
 \end{aligned} \tag{26}$$

for $x \in \partial\Omega$, where we have used (25). Therefore, by Theorem 7, the operator \mathcal{F} has at least one fixed point which in turn implies that the problem (2) has at least one solution. \square

Theorem 10. *Assume that $f : [0, T] \times \mathcal{R} \rightarrow \mathcal{R}$ is a continuous function satisfying the condition*

$$|f(t, x) - f(t, y)| \leq L|x - y|, \quad \forall t \in [0, T], \quad x, y \in \mathcal{R}, \tag{27}$$

with $L\kappa < 1$, where

$$\begin{aligned}
 \kappa = T^q \left[\frac{3}{2\Gamma(q+1)} + \frac{\Gamma(2-p)}{2\Gamma(q-p+1)} + \frac{\Gamma(2-p)}{4\Gamma(q-p)} \right. \\
 \left. \times \left\{ 1 - \frac{2\Gamma(2-p)}{\Gamma(3-p)} + 2 \left(\frac{\Gamma(2-p)}{\Gamma(3-p)} \right)^2 \right\} \right].
 \end{aligned} \tag{28}$$

Then the problem (2) has a unique solution.

Proof. Let us fix $\sup_{t \in [0, T]} |f(t, 0)| = M < \infty$ and select

$$r \geq \frac{M\kappa}{1 - L\kappa}, \tag{29}$$

where κ is given by (28). Then we show that $\mathcal{F}B_r \subset B_r$, where $B_r = \{x \in \mathfrak{C} : \|x\| \leq r\}$. For $x \in B_r$, we have

$$\begin{aligned}
 |(\mathcal{F}x)(t)| \\
 \leq \int_0^t \frac{(t-s)^{q-1}}{\Gamma(q)} |f(s, x(s)) - f(s, 0) + f(s, 0)| ds
 \end{aligned}$$

$$\begin{aligned}
 & + \frac{1}{2} \int_0^T \frac{(T-s)^{q-1}}{\Gamma(q)} |f(s, x(s)) - f(s, 0) + f(s, 0)| ds \\
 & + \frac{\Gamma(2-p)(T-2t)}{2T^{1-p}} \\
 & \times \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} |f(s, x(s)) - f(s, 0) + f(s, 0)| ds \\
 & + \Gamma(2-p)T^{p-1} \\
 & \times \left(T^2 - 2t^2 - \frac{2T^2\Gamma(2-p)}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right) (4)^{-1} \\
 & \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} |f(s, x(s)) - f(s, 0) + f(s, 0)| ds \\
 \leq & (Lr + M) \\
 & \times \left\{ \frac{2|t|^q + T^q}{2\Gamma(q+1)} + \frac{\Gamma(2-p)T^{q-1}|T-2t|}{2\Gamma(q-p+1)} \right. \\
 & \left. + \frac{\Gamma(2-p)T^{q-2}}{4\Gamma(q-p)} \right. \\
 & \left. \times \left| T^2 - 2t^2 - \frac{2\Gamma(2-p)T^2}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right| \right\} \\
 \leq & (Lr + M) \\
 & \times T^q \left[\frac{3}{2\Gamma(q+1)} + \frac{\Gamma(2-p)}{2\Gamma(q-p+1)} + \frac{\Gamma(2-p)}{4\Gamma(q-p)} \right. \\
 & \left. \times \left\{ 1 - \frac{2\Gamma(2-p)}{\Gamma(3-p)} + 2 \left(\frac{\Gamma(2-p)}{\Gamma(3-p)} \right)^2 \right\} \right] \\
 = & (Lr + M) \kappa \leq r.
 \end{aligned} \tag{30}$$

Thus we get $\mathcal{F}x \in B_r$. Now, for $x, y \in \mathfrak{C}$ and for each $t \in [0, T]$, we obtain

$$\begin{aligned}
 & |(\mathcal{F}x)(t) - (\mathcal{F}y)(t)| \\
 \leq & \left\{ \int_0^t \frac{(t-s)^{q-1}}{\Gamma(q)} \|f(s, x(s)) - f(s, y(s))\| ds \right. \\
 & + \frac{1}{2} \int_0^T \frac{(T-s)^{q-1}}{\Gamma(q)} \|f(s, x(s)) - f(s, y(s))\| ds \\
 & + \frac{\Gamma(2-p)(T-2t)}{2T^{1-p}} \\
 & \times \int_0^T \frac{(T-s)^{q-p-1}}{\Gamma(q-p)} \|f(s, x(s)) - f(s, y(s))\| ds \\
 & \left. + \Gamma(2-p)T^{p-1} \right\}
 \end{aligned}$$

$$\begin{aligned}
 & \times \left(T^2 - 2t^2 - \frac{2T^2\Gamma(2-p)}{\Gamma(3-p)} + \frac{4tT\Gamma(2-p)}{\Gamma(3-p)} \right) (4)^{-1} \\
 & \times \int_0^T \frac{(T-s)^{q-p-2}}{\Gamma(q-p-1)} \|f(s, x(s)) - f(s, y(s))\| ds \Big\} \\
 \leq & L \|x - y\| \\
 & \times T^q \left[\frac{3}{2\Gamma(q+1)} + \frac{\Gamma(2-p)}{2\Gamma(q-p+1)} + \frac{\Gamma(2-p)}{4\Gamma(q-p)} \right. \\
 & \left. \times \left\{ 1 - \frac{2\Gamma(2-p)}{\Gamma(3-p)} + 2 \left(\frac{\Gamma(2-p)}{\Gamma(3-p)} \right)^2 \right\} \right] \\
 = & \kappa L \|x - y\|,
 \end{aligned} \tag{31}$$

which, in view of the condition $\kappa L < 1$ (κ is given by (28)), implies that the operator \mathcal{F} is a contraction. Hence, by Banach's contraction mapping principle, the problem (2) has a unique solution. \square

Example 11. Consider the following antiperiodic fractional boundary value problem:

$$\begin{aligned}
 {}^c D^q x(t) &= (t^2 + 1) e^{-x^2(t)} \ln(4 + 3\sin^2 x(t)), \\
 & 0 < t < 1, \quad 2 < q \leq 3, \\
 x(0) &= -x(1), \\
 {}^c D^p x(0) &= -{}^c D^p x(1), \quad {}^c D^{p+1} x(0) = -{}^c D^{p+1} x(1), \\
 & 0 < p < 1.
 \end{aligned} \tag{32}$$

Clearly $|f(t, x(t))| \leq (3 \ln 7)$. So, the hypothesis of Theorem 8 holds. Therefore, the conclusion of Theorem 8 applies to antiperiodic fractional boundary value problem (32).

Example 12. Consider the following antiperiodic fractional boundary value problem:

$$\begin{aligned}
 {}^c D^{5/2} x(t) &= \frac{L}{2} (x(t) + \tan^{-1} x(t)) + \sqrt{1 + \sin^3 t}, \\
 & L > 0, \quad t \in [0, 2], \\
 x(0) &= -x(2) \\
 {}^c D^{3/4} x(0) &= -{}^c D^{3/4} x(2), \quad {}^c D^{7/4} x(0) = -{}^c D^{7/4} x(2),
 \end{aligned} \tag{33}$$

where $q = 5/2$, $p = 3/4$, $p + 1 = 7/4$ $f(t, x) = L(x + \tan^{-1} x)/2 + \sqrt{1 + \sin^3 t}$, and $T = 2$. Clearly,

$$\begin{aligned}
 |f(t, x) - f(t, \bar{x})| &\leq \frac{L}{2} (|x - \bar{x}| + |\tan^{-1} x - \tan^{-1} \bar{x}|) \\
 &\leq L(|x - \bar{x}|),
 \end{aligned} \tag{34}$$




where we have used the fact that $|(\tan^{-1}y)'| = 1/(1+y^2) < 1$. Further,

$$\begin{aligned} \kappa &= T^q \left[\frac{3}{2\Gamma(q+1)} + \frac{\Gamma(2-p)}{2\Gamma(q-p+1)} + \frac{\Gamma(2-p)}{4\Gamma(q-p)} \right. \\ &\quad \left. \times \left\{ 1 - \frac{2\Gamma(2-p)}{\Gamma(3-p)} + 2 \left(\frac{\Gamma(2-p)}{\Gamma(3-p)} \right)^2 \right\} \right] \quad (35) \\ &= \frac{16}{5} \sqrt{\frac{2}{\pi}} + \frac{526\Gamma(1/4)}{525\Gamma(3/4)}. \end{aligned}$$

With $L < 1/\kappa$, all the assumptions of Theorem 10 are satisfied. Hence, the fractional boundary value problem (33) has a unique solution on $[0, 2]$.

References

- [1] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives, Theory and Applications*, Gordon and Breach, Yverdon, Switzerland, 1993.
- [2] I. Podlubny, *Fractional Differential Equations*, vol. 198, Academic Press, San Diego, Calif, USA, 1999.
- [3] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, vol. 204 of *North-Holland Mathematics Studies*, Elsevier Science, Amsterdam, The Netherlands, 2006.
- [4] J. Sabatier, O. P. Agrawal, and J. A. T. Machado, Eds., *Advances in Fractional Calculus: Theoretical Developments and Applications in Physics and Engineering*, Springer, Dordrecht, The Netherlands, 2007.
- [5] D. Baleanu, K. Diethelm, E. Scalas, and J. J. Trujillo, *Fractional Calculus Models and Numerical Methods. Series on Complexity, Nonlinearity and Chaos*, vol. 3, World Scientific, Boston, Mass, USA, 2012.
- [6] Z. Bai, "On positive solutions of a nonlocal fractional boundary value problem," *Nonlinear Analysis: Theory, Methods and Applications A*, vol. 72, no. 2, pp. 916–924, 2010.
- [7] V. Gafiychuk and B. Datsko, "Mathematical modeling of different types of instabilities in time fractional reaction-diffusion systems," *Computers and Mathematics with Applications*, vol. 59, no. 3, pp. 1101–1107, 2010.
- [8] D. Băleanu and O. G. Mustafa, "On the global existence of solutions to a class of fractional differential equations," *Computers and Mathematics with Applications*, vol. 59, no. 5, pp. 1835–1841, 2010.
- [9] D. Băleanu, O. G. Mustafa, and D. O'Regan, "A Nagumo-like uniqueness theorem for fractional differential equations," *Journal of Physics A*, vol. 44, no. 39, Article ID 392003, 6 pages, 2011.
- [10] B. Ahmad and J. J. Nieto, "Riemann-Liouville fractional integro-differential equations with fractional nonlocal integral boundary conditions," *Boundary Value Problems*, vol. 2011, no. 36, 9 pages, 2011.
- [11] B. Ahmad and S. K. Ntouyas, "A four-point nonlocal integral boundary value problem for fractional differential equations of arbitrary order," *Electronic Journal of Qualitative Theory of Differential Equations*, vol. 2011, no. 22, 15 pages, 2011.
- [12] N. J. Ford and M. L. Morgado, "Fractional boundary value problems: analysis and numerical methods," *Fractional Calculus and Applied Analysis*, vol. 14, no. 4, pp. 554–567, 2011.
- [13] A. Aghajani, Y. Jalilian, and J. J. Trujillo, "On the existence of solutions of fractional integro-differential equations," *Fractional Calculus and Applied Analysis*, vol. 15, no. 1, pp. 44–69, 2012.
- [14] B. Ahmad and S. K. Ntouyas, "A note on fractional differential equations with fractional separated boundary conditions," *Abstract and Applied Analysis*, vol. 2012, Article ID 818703, 11 pages, 2012.
- [15] B. Datsko and V. Gafiychuk, "Complex nonlinear dynamics in subdiffusive activator-inhibitor systems," *Communications in Nonlinear Science and Numerical Simulation*, vol. 17, no. 4, pp. 1673–1680, 2012.
- [16] B. Ahmad, J. J. Nieto, A. Alsaedi, and M. El-Shahed, "A study of nonlinear Langevin equation involving two fractional orders in different intervals," *Nonlinear Analysis: Real World Applications*, vol. 13, no. 2, pp. 599–606, 2012.
- [17] B. Ahmad and J. J. Nieto, "Existence of solutions for anti-periodic boundary value problems involving fractional differential equations via Leray-Schauder degree theory," *Topological Methods in Nonlinear Analysis*, vol. 35, no. 2, pp. 295–304, 2010.
- [18] B. Ahmad, "Existence of solutions for fractional differential equations of order $q \in (2,3]$ with anti-periodic boundary conditions," *Journal of Applied Mathematics and Computing*, vol. 34, no. 1-2, pp. 385–391, 2010.
- [19] R. P. Agarwal and B. Ahmad, "Existence of solutions for impulsive anti-periodic boundary value problems of fractional semi-linear evolution equations," *Dynamics of Continuous, Discrete and Impulsive Systems A*, vol. 18, no. 4, pp. 457–470, 2011.
- [20] B. Ahmad, "New results for boundary value problems of nonlinear fractional differential equations with non-separated boundary conditions," *Acta Mathematica Vietnamica*, vol. 36, no. 3, pp. 659–668, 2011.
- [21] G. Wang, B. Ahmad, and L. Zhang, "Impulsive anti-periodic boundary value problem for nonlinear differential equations of fractional order," *Nonlinear Analysis: Theory, Methods and Applications A*, vol. 74, no. 3, pp. 792–804, 2011.
- [22] B. Ahmad and J. J. Nieto, "Anti-periodic fractional boundary value problems," *Computers and Mathematics with Applications*, vol. 62, no. 3, pp. 1150–1156, 2011.
- [23] Y. Chen, J. J. Nieto, and D. O'Regan, "Anti-periodic solutions for evolution equations associated with maximal monotone mappings," *Applied Mathematics Letters*, vol. 24, no. 3, pp. 302–307, 2011.
- [24] B. Ahmad and J. J. Nieto, "Anti-periodic fractional boundary value problems with nonlinear term depending on lower order derivative," *Fractional Calculus and Applied Analysis*, vol. 15, no. 3, pp. 451–462, 2012.
- [25] D. R. Smart, *Fixed Point Theorems*, Cambridge University Press, Cambridge, UK, 1974.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

