

Chrysymenia wrightii (Rhodymeniales, Rhodophyta) - a new non-native species for the European Atlantic Coast

Ignacio Bárbara^{1*}, Sook-Young Lee², Viviana Peña¹, Pilar Díaz¹, Javier Cremades¹, Jung Hyun Oak³ and Han-Gu Choi²

¹Facultad de Ciencias, Universidad de A Coruña, Campus de la Zapateira s/n, 15071, Spain

²Korea Polar Research Institute, KORDI, 7-50 Songdo-dong, Incheon 406-840, Korea

³Marine Research Institute, Pusan National University, Busan 609-735, Korea

*Corresponding author

E-mail: barbara@udc.es

Received 3 July 2008; accepted in revised form 15 September 2008; published online 18 December 2008

Abstract

Chrysymenia wrightii, originally described from Japan, was found for the first time from the European Atlantic coast. It was collected in several subtidal rocky bottom habitats (9-14 meters depth) of the Ría de Arousa (Galicia, NW Spain). In this work, a description of Galician gametophytic and tetrasporophytic plants is provided. They are similar to the Mediterranean and Japanese plants. DNA sequence data of materials from Galicia and Korea were investigated using nuclear SSU and ITS1-5.8S-ITS2 rDNA and plastid *rbcL* sequences. No genetic variation was observed in the SSU, and only one substitution was detected in ITS and *rbcL* data between Galician and Korean samples, respectively. Our molecular data indicate that the Galician populations of *C. wrightii* are probably due to a recent introduction event from the northwest Pacific. Despite the fact that *C. wrightii* was formerly recorded as a new non-native species from a Mediterranean hotspot (Thau Lagoon, 1987-1989), it was never reported for the European Atlantic coast. After 30 years of its first report in the Mediterranean Sea, the simultaneous occurrence of *C. wrightii* in four subtidal localities of NW Spain suggests that this species may have colonized the region unnoticed for several years which may be related to its subtidal habitat and short life cycle. Similarly to the Mediterranean Sea, the introduction of *C. wrightii* on the European Atlantic coasts could be enhanced by the intensive aquaculture widely spread along the Galician coast. Curiously, *C. wrightii* is apparently restricted to both areas, the Ría de Arousa and the Thau Lagoon. In addition, many non-native species markedly represented in Galicia since the 1980's were firstly reported in the Ría de Arousa. Consequently, the Ría de Arousa should be considered an outstanding European Atlantic hotspot of introduced marine species similar to the Solent region (South of England) or the Thau Lagoon.

Key words: *Chrysymenia wrightii*, Spain, non-native species, Atlantic, seaweed, aquaculture

Introduction

The introduction of non-native species is mainly caused by vessels and aquaculture (Streftaris et al. 2005). In the Galician region (NW Spain), fishing and aquaculture industry have a great development. They have favoured the interchange of biological material, which is

responsible for changes in the natural communities. During the last years, several non-native species are being locally abundant in the Galician rías. *Sargassum muticum* (Yendo) Fensholt, *Undaria pinnatifida* (Harvey) Suringar, *Lomentaria hakodatensis* Yendo, *Grateloupia turuturu* Yamada, *G. subpectinata* Holmes, *Gracilaria vermiculophylla* (Ohmi) Papenfuss, *Heterosiphonia japonica* Yendo, *Dasya sessilis*

Yamada, *Neosiphonia harveyi* (J. Bailey) M.S. Kim, H.-G. Choi, Guiry & G.W. Saunders and *Ulva pertusa* Kjellman have been reported along the Galician coasts, and most of them are widely spread along the coast and the Galician rías (Pérez-Cirera et al. 1989; Santiago Caamaño 1990; López Rodríguez et al. 1991; ICES 1992; Cremades Ugarte 1995; Bárbara and Cremades 1996; 2004, Veiga et al. 1998; Pérez-Ruzafa et al. 2002; Rueness 2005; Bárbara et al. 2002, 2003, 2005; De Clerck et al. 2005; Barreiro et al. 2006; Cremades Ugarte et al. 2006; Freire et al. 2006; Peña and Bárbara 2006; Baamonde López et al. 2007 and López Figueroa et al. 2007).

Although *Chrysymenia wrightii* (Harvey) Yamada was reported as non-native seaweed for the European coast (Thau Lagoon, Mediterranean Sea) (Ben Maïz et al. 1987), it has not been detected on the European Atlantic coast. In the NW of Iberian Peninsula, no *Chrysymenia* species are known. Solely, *C. ventricosa* was reported in the Cantabric Sea (Gorostiaga et al. 2004; Llera González and Álvarez Raboso 2007) whereas the rest of the Iberian records for this warm-temperate species are known from the South of Spain (Conde et al. 1996) and the Mediterranean Sea (Ballesteros 1989). During current surveys in subtidal maërl beds in the Galician rías, several specimens of the northwest Pacific species *C. wrightii* were collected from rocky bottom areas. The aim of this work is to add *C. wrightii* as a new non-native species to the European Atlantic seaweeds catalogue and describe its morphology, habitat, current distribution and extent of its populations, as well as the genetic variation of the Atlantic and Pacific populations.

Material and Methods

Plants were first collected (October 2005) by SCUBA diving in a subtidal rocky bottom habitat covered with mud (9-12 meters depth) of the Ría de Arousa during the explorations of the Galician maërl beds. From a new subtidal exploration along this ría (September and October 2007) we found *C. wrightii* in the original site examined in 2005 and in three new localities, Cambados, Isote Galíñeiro and Tragove (Figure 1, Annex 1) which are located 11 km from the first locality. Samples were preserved in 4% formalin seawater at 4°C in darkness. In order to study reproductive structures, sterile plants collected in Cabo Cruz (September 2007) were



Figure 1. *Chrysymenia wrightii* in Galicia (NW Spain), localization of the non-native plants along the Ría de Arousa (Galicia, Spain)

cultured in sterile seawater (31-32 psu) at 16°C and 4:20 light:dark photoperiod. Plants were dissected by hand with a stainless steel razor blade. Herbarium specimens were deposited at the herbarium of the University of Santiago de Compostela (SANT-Algae).

In order to study the genetic variation between Galician and Korean individuals of *C. wrightii*, several silica-dried specimens from both areas (Annex 2) were compared at the Polar Biodiversity Laboratory, Korea Polar Research Institute (KOPRI). Genomic DNA from samples was extracted using DNeasy® Plant Mini Kit (Qiagen, Hilden, Germany). After DNA extraction, the rest of silica-dried samples were deposited at the KOPRI herbarium. The nuclear SSU and ITS rDNA, and plastid *rbcL* were amplified from total genomic DNA using polymerase chain reaction (PCR) and the primer combinations of

Saunders and Kraft (1994, 1996) for SSU, Tai et al. (2001) for ITS1-5.8S-ITS2, and Freshwater and Rueness (1994) for *rbcL*. Agarose gel-purification or direct purification with High Pure™ PCR Product Purification Kit (Roche Diagnostics, Indianapolis, IN, USA) or LaboPass PCR Purification Kit (COSMO Genetech, Seoul, Korea) was used to clean PCR products. DNA purified using this method was sequenced using the BigDye™ terminator cycle sequencing ready reaction kit (PE Applied Biosystems [ABI], Foster City, CA, USA). Sequence data were collected using an ABI PRISM 3730 DNA Analyzer, and were edited using the SeqEd DNA sequence editor (ABI) software package. The edited sequences were aligned relative to one another using the SeqPup multiple alignment program (Gilbert 1995) and MacClade 4 program (Maddison and Maddison 2003).

The final alignment for SSU, *rbcL* and ITS1-5.8S-ITS2 consisted of three, three and seven taxa (Annex 2), respectively. The sequences of

the 1771, 1526 and 703 aligned nucleotide positions of SSU, *rbcL* and ITS1-5.8S-ITS2 data were edited to remove the 5' and 3' PCR primer regions (Saunders and Kraft 1994; Freshwater and Rueness 1994; Tai et al. 2001), as well as ambiguously aligned regions, to yield 1723, 1448 and 658 base pairs for phylogenetic inference, respectively. Pairwise distance for SSU, *rbcL* and ITS1-5.8S-ITS2 data were performed using PAUP* 4.0b10 for Macintosh (Swofford 2002).

Results and Discussion

Galician plants of *Chrysiomenia wrightii* are similar to the ones described for the Mediterranean coast (Ben Maiz et al. 1987) and the Japanese coast (Yamada 1932; Lee 1978). They are 15-40 cm high, red-brownish in colour, terete and hollow (Figure 2). The thallus is gelatinous and tender, attached by a discoid holdfast and




Figure 2. *Chrysiomenia wrightii* in Galicia (NW Spain), habit. A. Tetrasporophytic plant (Cabo Cruz, October 2005). B. Gametophytic plant with cystocarps (Islote Galiñeiro, October 2007). C. Herbarium material from Cambados, 7-X-2007 (SANT-Algae 19525). D. Apical branches, constricted at the base and tapering at the tips. Scale bar A-C = 5 cm, D = 5 mm (Photographs by Ignacio Bárbara)




Figure 3. *Chrysymenia wrightii* in Galicia, vegetative structure. A. Transverse section of an apical branch showing the hollow thallus. B-C. Cortical, subcortical and hyphaelike filaments in transverse section. D. Cortical cells in surface view, outer part. E. Hyphaelike filaments and gland cells in surface view, inner part. F. Gland cells. G. Hyphaelike filaments. Scale bar = 200 μm (Photographs by Ignacio Bárbara)

cylindrical axes, 3-4 mm at the base to 1-2 mm at the apex. Plants grow monopodially, 2-4 times branched (alternate, opposite or irregular) with branches constricted at the base and tapering at the tips. The cortical layer consists of 2-3 small cells (5-8 x 6-12 μm) and the subcortical layer has 3-5 large cells and inner irregular rizoidal cells (Figure 3). The medullary layer of 3-5 cells is loosely and irregularly arranged. Medullary cells are hyaline, elongated (130-400 x 60-90 μm). Gland cells (up to 70 μm diameter) and hyphae-like filaments (2-5 cells) are developed from innermost medullary cells, solitary or in aggregation (Figure 3) as it is commented by Yamada (1932).

Gametophytic plants are covered by numerous cystocarps along the main axe and 1-2 order branches. Mature cystocarps are subspherical (460-950 (1200) μm in diameter) with a carpostome (80-120 μm in diameter) and carpospores (17-20 x 20-30 μm in diameter) (Figure 4). Cystocarps match with the descriptions of Lee (1978) and Yamada (1932). Whereas, Ben Maiz et al. (1987) described cystocarps (500-800 μm in diam.) without ostiole for the Mediterranean plants. In Galicia, this feature is observed only in immature cystocarps. Male plants were not observed. Tetrasporangia divided cruciately (28-42 x 25-30 μm) scattered in the cortical layer and originate from inner cortical cells (Figure 4).




Figure 4. *Chrysymenia wrightii* in Galicia, reproductive structure. A. Tetrasporangial plant in transverse section. B. Tetrasporangia under cortical cells, surface view. C. Cystocarp subspherical with carpostome. D. Mature spherical carpospores. E. Adventitious short branches from cystocarps (arrows). Scale bar A, C-D = 200 μ m, B = 50 μ m, E = 2 cm (Photographs by Ignacio Bárbara)

Galician plants show a morphological variability in size and diameter of axes. Plants are profuse in branching, second order branches being abundant. In general, gametophytic plants are more irregular in form and branching compared to the tetrasporophytic plants, because adventitious short branches (similar in form to apical branches) grow from the pericarp of senescent cystocarps. This growth process is quite peculiar since reproductive structures are usually the last growth stadium, but this species apparently have a high cell activity. Profusion in branching is a good strategy for non-native and invasive species providing lots of vegetative propagules by fragmentation. Although we have not observed new individuals of *C. wrightii* growing from the cystocarpic branches, they could provide an excellent way of spreading along the Atlantic coast of Europe as it was pointed out for other non-native species with effective dispersal by vegetative propagation

(Eno et al. 1997; Bjaerke and Rueness 2004; Husa and Sjötnun 2006).

Three SSU rDNA, three *rbcL* and seven ITS1-5.8S-ITS2 rDNA sequences were completed from seven samples of two different localities in Galicia and three different sites in Korea and were deposited in GenBank (Annex 2). No ambiguities were observed in the sequence data. Comparisons with sequences retrieved from GenBank showed that the SSU rDNA sequences were identical with that of *C. wrightii* from Japan (Saunders et al. 1999; AF117129). All sequence data for two samples (CH1407 and CH1536) from Galicia were identical to each other. No genetic variation was seen in the SSU, and only one substitution was observed in *rbcL* (T \leftrightarrow C in position 914; data not shown) and ITS data (A \leftrightarrow G in position 423; Figure 5) between Galician and all Korean samples, respectively. In addition a deletion (T in position 169) and an insertion (T in position 538) were also observed

in the ITS1 and ITS2 region, respectively between Galician and four Korean ones (CH1727, CH1728, CH1729 and CH1774; Figure 5). Only one deletion (T in position 169) was observed between the sample from Samchuk (CH1770) and the other four samples from Korea (Figure 5). Our molecular data indicate that the *C. wrightii* populations in Galicia are probably due to a recent introduction event from

somewhere of the northwest Pacific near Korean coast.

In the Mediterranean Sea the vector of introduction for *Chrysomenia wrightii* is likely to be mollusc culture (Verlaque 2001; Cormaci et al. 2004). Taking into account that intensive aquaculture is widely spread on the Galician coast, the introduction of *C. wrightii* was likely enhanced by this vector. In the NW Spain,




Figure 5. A sub-alignment (positions 101 to 200 and 401 to 600) of ITS1-5.8S-ITS2 rDNA sequences for five Korean (Chr.H1727, Chr.H1728, Chr.H1729, Jangseungpo; Chr.H1774, Yangyang; Chr.H1770, Samchuk) and two Galician samples (Chr.H1407, Boiro; Chr.H1536, Islote Galiñeiro). Bold characters (positions 169, 423 and 538 indicate a deletion, a substitution (G→A) and an insertion between Korean and Galician samples, respectively). A dash (-) indicates a gap in the multiple alignment; a dot (.) indicates that a taxon has the same character state for that site as in Chr.H1727

aquaculture has played an important role in the introduction of other non-native species especially since the 1980's. Examples include *Sargassum muticum*, *Undaria pinnatifida*, *Lomentaria hakodatensis*, *Grateloupia turuturu*, *G. subpectinata*, *Gracilaria vermiculophylla*, *Heterosiphonia japonica*, *Dasya sessilis* and *Ulva pertusa*.

Chrysomenia wrightii was added as a new non-native species for the European coast (Thau Lagoon, Mediterranean Sea), based on collections from 1978 to 1985 (Ben Maïz et al. 1987). Subsequently, the species is regularly found in the same area (which is a hotspot of introduced species) (Verlaque 2001, Verlaque et al. 2007) Many other non-native Japanese species such as *Saccharina japonica* (J.E. Areschoug) C.E. Lane, C. Mayes, Druehl & G.W. Saunders, *Ahnfeltiopsis flabelliformis* (Harvey) Masuda and *Sphaerotrachia firma* (E. Gepp) Zinova are equally restricted to the Thau Lagoon with

earlier introductions from 1971 to 1988 (Verlaque 2001, Verlaque et al. 2007), although, *S. japonica* has not been detected in the Thau Lagoon since 1989 (Verlaque 2001). In Galicia, *Grateloupia subpectinata* Holmes is an example of a non-native species which has not expanded along the coast. Hitherto, it was only known in 6 localities of the Ría de Arousa (López Rodríguez et al. 1991; Bárbara et al. 2002) and since the 1990's it has not colonized new localities in other Galician rías. At the present, *C. wrightii* is restricted to both the Mediterranean coast (Thau Lagoon) and the Atlantic coast (4 localities in the Ría de Arousa). Taking into account that the Ría de Arousa provides the new record of *C. wrightii* on the Atlantic coasts, as it happened with other non-native species which are now widely distributed, we conclude that the Ría de Arousa should be considered as an interesting European Atlantic hotspot for the introduction of marine species such as the Solent region in

England (Farnham 1980) and the Thau Lagoon in the Mediterranean France (Verlaque 2001).

The present record of *Chrysymenia wrightii* in the Ría de Arousa occurred 30 years after its detection in the Mediterranean. The subtidal habitat (9-14 meters depth) where the Galician populations were observed could have favoured an unnoticed occurrence for several years. In contrast in the Mediterranean Sea it could have been detected easier because it occurs at shallow localities, from 0.5 to 6 m depth (Ben Maïz et al. 1987). The short life cycle of *C. wrightii* could further complicate its detection. On the Japanese coasts it is described as an annual species (Lee 1978) which appears for a short period, May to October, whereas in the Mediterranean Sea it was found from January to November (Ben Maïz et al. 1987). However, in Galicia it was only collected in September and October despite many subtidal explorations (2006-2007) carried out along the Ría de Arousa in different months, habitats and water depths. Based on both features that complicate the detection of *C. wrightii* in Galicia (subtidal habitat and short life cycle), it would be possible that *C. wrightii* has been introduced earlier in the Ría de Arousa, 1-2 decades ago or more, but it has been unnoticed. Arguably, *C. wrightii* could also occur unnoticed nowadays along other Atlantic coasts which may only be proven with further subtidal surveys. New studies could also verify if the present record is an isolated case of introduction that will not thrive along the Atlantic Iberian Peninsula or if it has already become an established member of the Galician non-native seaweed flora or even if *C. wrightii* could be considered a future invasive species. Currently *C. wrightii* occurs in two areas of the Ría de Arousa: (a) locality 1 (1 ha surface) and (b) localities 2-4 (100 ha potential surface). Both are located in the vicinity of aquaculture sites where non-native species like *Undaria pinnatifida* or *Grateloupia turuturu* were firstly recorded for the Spanish coast.

Historical reports and our molecular data indicate that the *C. wrightii* populations in Galicia were recently introduced and originate from somewhere in the northwest Pacific and may have used the Mediterranean coast as a stepping stone. To complete the information about this non-native species, extensive samplings of plants and more variable genetic markers could more clearly locate the source populations and further identify the most likely dispersal vector(s).

Acknowledgements

We thank Dr. Marc Verlaque for verifying the identification of the first Galician plants (October 2005) and his revision of the manuscript. We thank to Carmen Lema for helping in the last field work. We are also grateful to Dr. Hyung-Seop Kim for kindly providing some samples. Contribution to the projects PGIDIT03PXIB10301PR (Xunta de Galicia) and CGL2006-03576/BOS (Ministerio de Educación y Ciencia, FEDER). This work was supported by a grant from the Basic Research Program of the Korea Science and Engineering Foundation (R01-2006-000-10312-0) and a grant from the Basic Research Program of the Korea Polar Research Institute project PE08060.

References

- Baamonde López S, Baspino Fernández I, Barreiro Lozano R, Cremades Ugarte J (2007) *Ulva pertusa* (Ulvales, Chlorophyta), a new alien seaweed on the European Atlantic coast. *Botanica Marina* 50: 267-274.
- Ballesteros E (1989) Checklist of benthic marine algae from Catalonia (North Western Mediterranean). *Treballs de l'Institut Botànic de Barcelona* 13:1-52
- Bárbara I and Cremades J (1996) Seaweeds of the Ría de A Coruña (NW Iberian Peninsula, Spain) *Botanica Marina*, 39: 371-388
- Bárbara I, Cremades J (2004) *Grateloupia lanceola* versus *Grateloupia turuturu* (Gigartinales, Rhodophyta) en la Península Ibérica. *Anales del Jardín Botánico de Madrid* 61(2): 103-118
- Bárbara I, Cremades J, Veiga AJ, López-Varela C, Dosil J, Calvo S, Peña V, López-Rodríguez MC (2002) Fragmenta Chorologica Occidentalia, Algae, 7814-7892. *Anales del Jardín Botánico de Madrid* 59(2): 292-297
- Bárbara I, Cremades J, Veiga AJ, López Rodríguez MC (2003) *Dasysiphonia* sp. (Ceramiales, Dasyaceae) nuevo rodófito alóctono para la Península Ibérica. *Anales del Jardín Botánico de Madrid* 60(2): 440-443
- Bárbara I, Cremades J, Calvo S, López Rodríguez MC, Dosil J (2005) Checklist of the benthic marine and brackish Galician algae (NW Spain). *Anales del Jardín Botánico de Madrid* 62(1): 69-100
- Barreiro R, Quintela M, Bárbara I, Cremades J (2006) RAPD differentiation between and invasive and a native species of *Grateloupia* (Rhodophyta) in Galicia (NW Spain). *Phycologia* 46: 213-217
- Ben Maïz N, Boudouresque CF, Riouall R, Lauret M (1987) Flore algale de l'Etang de Thau (France, Méditerranée): sur la présence d'une Rhodymeniale d'origine japonaise, *Chrysymenia wrightii* (Rhodophyta). *Botanica Marina* 30(5): 357-364
- Bjaerke MR, Rueness EK (2004) Effects of temperature and salinity on growth, reproduction and survival in the introduced red alga *Heterosiphonia japonica* (Ceramiales, Rhodophyta). *Botanica Marina* 47: 373-380
- Conde F, Flores-Moya A, Soto J, Altamirano M, Sánchez A (1996) Check-list of Andalusia (S. Spain) seaweeds. III. Rhodophyceae. *Acta Botanica Malacitana* 21: 7-33

- Cormaci M, Furnari, G, Giaccone G, Serio D (2004) Alien Macrophytes in the Mediterranean Sea: A review. Recent research Development Environmental Biology 1: 153-202
- Cremades Ugarte J (1995) A introducción de algas marinha alóctonas nas costas de Galiza. Cerna 16: 12-15
- Cremades Ugarte J, Freire Gago Ó, Peteiro García C (2006) Biología, distribución e integración del alga alóctona *Undaria pinnatifida* (Laminariales, Phaeo-phyta) en las comunidades bentónicas de las costas de Galicia (N.O. Península Ibérica). Anales del Jardín Botánico de Madrid 63(2): 169-187
- De Clerck O, Gavio B, Fredericq S, Bárbara I, Coppejans E (2005) Systematics of *Grateloupia filicina* (Halymeniaceae, Rhodophyta), based on *rbcL* sequence analyses and morphological evidence, including the reinstatement of *G. minima* and the description of *G. capensis* spec. nov. Journal of Phycology 41: 391-410
- Eno NC, Clark RA, Sanderson WG (1997) Non-native marine species in British waters: a review and directory. Joint Nature Conservation Committee, Peterborough, 83 pp.
- Farnham WF (1980) Studies on aliens in the marine flora of Southern England. In: The Shore Environment: Ecosystems, vol. 2. Ecosystems. Ed. By Price JH, Irvine DEG and Farnham WF. Academic Press, London, pp 875-914
- Freire Ó, Peteiro C, Cremades J (2006) La integración del alga alóctona *Undaria pinnatifida* (Laminariales, Phaeophyceae) en la flora de las costas atlánticas peninsulares. Algas, Boletín de la Sociedad Española de Ficológia 35: 25-32
- Freshwater DW, Rueness J (1994) Phylogenetic relationships of some European *Gelidium* (Gelidiales, Rhodophyta) species, based on *rbcL* nucleotide sequence analysis. Phycologia 33: 187-194
- Gilbert DG (1995) SeqPup, a biosequence editor and analysis application. Biological Department, Indiana University, Bloomington
- Gorostiaga JM, Santolaria A, Secilla A, Casares C, Díez I (2004) Check-list of the Basque Coast Benthic Algae (N Spain). Anales del Jardín Botánico de Madrid 61: 155-180
- Husa V, Sjötn K (2006) Vegetative reproduction in "*Heterosiphonia japonica*" (Dasyaceae, Ceramiales, Rhodophyta), an introduced red alga on European coasts. Botanica Marina 49: 191-199
- ICES (1992). Report of the Working Group on Introductions and Transfer of Marine Organisms 1992/7. International Council for the exploration of the Sea (ICES). Lisbon
- Lee IK (1978) Studies on Rhodymeniales from Hokkaido. Journal of the Faculty of Sciences, Hokkaido University, Serie V (Botany) 11: 1-194
- Llera González EM, Álvarez Raboso J (2007) Algas Marinas de Asturias. Consejería de Medio Ambiente, Ordenación del Territorio e Infraestructuras. Oviedo
- López Figueroa F, Korbee N, De Clerk O, Bárbara I, Argall E (2007) Characterization of *Grateloupia lanceola* (Halymeniaceae, Rhodophyta), an obscure foliose *Grateloupia* from the Iberian Peninsula, based on morphology, *rbcL* sequences and mycosporine-like amino acid (MAA) composition. European Journal of Phycology 42(3): 231-242
- López Rodríguez MC, Cremades J, Bárbara I (1991) Fragmenta Chorologica Occidentalia, Algae, 3260-3284. Anales del Jardín Botánico de Madrid 49(1): 97-100
- Maddison W, Maddison D (2003) MacClade, 4.06. Sinauer Associates, Sunderland, MA, USA
- Peña V, Bárbara I (2006) Revision of the genus *Dasya* (Ceramiales, Rhodophyta) in Galicia (NW Spain) and the addition of a new alien species *Dasya sessilis* Yamada for the European Atlantic coasts. Anales del Jardín Botánico de Madrid 63(1): 13-26
- Pérez-Cirera JL, Cremades J, Bárbara I (1989) Precisiones sistemáticas y sinecológicas sobre algunas algas nuevas para Galicia o para las costas atlánticas de la Península Ibérica. Anales del Jardín Botánico de Madrid 46(1): 35-45
- Pérez-Ruzafa I, Menéndez JL, Salinas JM (2002) Mapas de distribución de algas marinas de la Península Ibérica e Islas Baleares. XV. *Undaria pinnatifida* (Harvey) Suringar (Laminariales, Fucophyceae). Botanica Complutensis 26: 147-151
- Rueness J (2005) Life history and molecular sequences of *Gracilaria vermiculophylla* (Gracilariales, Rhodophyta), a new introduction to European waters. Phycologia 44: 120-128
- Santiago Caamaño J, Durán Neira C, Acuña Castroviejo R (1990) Aparición de *Undaria pinnatifida* en las costas de Galicia (España). Un nuevo caso en la problemática de introducción de especies foráneas. Informes técnicos del Centro de Investigaciones Submarinas 3: 1-45
- Saunders GW, Kraft GT (1994) Small-subunit rRNA gene sequences from representatives of selected families of the Gigartinales and Rhodymeniales (Rhodophyta) I. Evidence for the Plocamiales ord. nov. Canadian Journal of Botany 72: 1250-1263
- Saunders GW, Kraft GT (1996) Small-subunit rRNA gene sequences from representatives of selected families of the Gigartinales and Rhodymeniales (Rhodophyta). 2. Recognition of the Halymeniaceae ord. nov. Canadian Journal of Botany 74: 694-707
- Saunders GW, Strachan IM, Kraft GT (1999) The families of the order Rhodymeniales (Rhodophyta): a molecular-systematic investigation with a description of Faucheaceae fam. nov. Phycologia 38: 23-40
- Strefitaris N, Zenetos A, Papanathanassiou E (2005) Globalisation in marine ecosystems: the story of non-indigenous marine species across European seas. Oceanography and Marine Biology, Annual Review 43: 419-453
- Swofford DL (2002) PAUP*: Phylogenetic analysis using parsimony (and other methods), version 4.0b10. Sinauer Associated, Inc., Sunderland, Massachusetts
- Tai V, Lindstrom SC, Saunders GW (2001) Phylogeny of the Dumontiaceae (Gigartinales, Rhodophyta) and associated families based on SSU rDNA and internal transcribed spacer sequence data. Journal of Phycology 37: 184-196
- Veiga AJ, Cremades J, Bárbara I (1998) Fragmenta Chorologica Occidentalia, Algae, 6283-6307. Anales del Jardín Botánico de Madrid 56(1): 121-123
- Verlaque M (2001) Checklist of the macroalgae of Thau Lagoon (Hérault, France) a hot spot of marine species introduction in Europe. Oceanologica Acta 24(1): 29-49
- Verlaque M, Boudouresque CF, Mineur F (2007). Oyster transfers as a vector for marine species introductions: a realistic approach based on the macrophytes. CIESM Workshop Monographs, Monaco, 32, pp 39-48
- Yamada Y (1932) Notes on some Japanese algae III. Journal of the Faculty of Sciences, Hokkaido University, Serie V (Botany), 1: 109-123

Annex 1

Galician samples of *Chrysymenia wrightii*

Map reference	Location	Geographic coordinates		Collection date	Depth	Substrate	Observations	Collector
		Latitude, N	Longitude, W					
1	Ría de Arousa, Cabo Cruz	42°36'32"	008°53'19"	3-X-2005	9-12 m	rocky	Tetrasporophytic plants living together to <i>Bonnemaisonia clavata</i> Hamel and <i>Scinaia interrupta</i> (A. DC.) Wynne	I. Bárbara, P. Díaz and V. Peña
				11-IX-2007	12 m	rocky with mud	Gametophytic plants	I. Bárbara, V. Peña and C. Lema
2	Ría de Arousa, Cambados	42°30'39"	008°50'47"	7-X-2007	14 m	rocky with mud	Sterile plants, living together to <i>Dasya sessilis</i> Yamada	I. Bárbara, V. Peña and C. Lema
3	Ría de Arousa, Islote Galiñeiro	42°31'02"	008°51'44"	7-X-2007	8 m	rocky with mud	Gametophytic and tetrasporophytic plants	I. Bárbara, V. Peña and C. Lema
4	Ría de Arousa, Travoge	42°30'58"	008°50'35"	7-X-2007	12 m	rocky with mud	Sterile plants.	I. Bárbara, C. Lema and V. Peña

Annex 2

Collection information for the samples used in the molecular study

Samples	Collection Details	GenBank		
		SSU	<i>rbcL</i>	ITS
CH1727	-2 m, Jangseungpo, southern coast of Korea, 23-II-2008, J. H. Oak	-	-	EU916720
CH1728	-2 m, Jangseungpo, southern coast of Korea, 23-II-2008, J. H. Oak	EU916714	EU916717	EU916721
CH1729	-2 m, Jangseungpo, southern coast of Korea, 23-II-2008, J. H. Oak	-	-	EU916722
CH1774	Intertidal, Yangyang, eastern coast of Korea, 24-V-2007, S.-M. Kim, C.-J. Kwon and H.-S. Kim	-	-	EU916724
CH1770	Intertidal, Samchuk, eastern coast of Korea, 9-III-2006, S.-M. Kim, C.-J. Kwon and H.-S. Kim	-	-	EU916723
CH1407	SANT-Algae 19058, Boiro, Cabo Cruz, Ría de Arousa, Spain, 3-X-2005, tetrasporophytic plant, I. Bárbara, P. Díaz and V. Peña	EU916712	EU916715	EU916718
CH1536	SANT-Algae 19514, Islote Galiñeiro, Ría de Arousa, Spain, 7-X-2007, tetrasporophytic plant, I. Bárbara, V. Peña and C. Lema	EU916713	EU916716	EU916719