

# Characterization of pathogenic vibrios isolated from bivalve hatcheries in Galicia, NW Atlantic coast of Spain. Description of *Vibrio tubiashii* subsp. *europaensis* subsp. nov

Susana Prado<sup>\*,1</sup>, Javier Dubert<sup>1</sup>, Juan L. Barja

Departamento de Microbiología y Parasitología, CIBUS—Facultad de Biología, Universidad de Santiago de Compostela, Santiago de Compostela, 15782, Spain

## ARTICLE INFO

DOI: [10.1016/j.syapm.2014.11.005](https://doi.org/10.1016/j.syapm.2014.11.005)

### Keywords:

*Vibrio tubiashii* subsp. *europaensis* subsp. nov.

Orientalis clade

MLSA

Bivalve hatchery

Larval pathogen

## ABSTRACT

The taxonomic position of the bivalve pathogen PP-638 was studied together with five similar isolates. The strains were isolated from flat oyster (*Ostrea edulis*) and Manila clam (*Venerupis philippinarum*) cultures during outbreaks of disease in two shellfish hatcheries (Galicia, NW Spain). The pathogenicity, previously established for PP-638, was demonstrated with all isolates and for several bivalve species, including the original hosts. On the basis of phenotypic characterization and 16S rRNA gene sequences, a tight group was defined within the genus *Vibrio*. Multilocus sequence analysis (MLSA) based on concatenated sequences of the 16S rRNA gene and the five housekeeping genes *recA*, *rpoA*, *pyrH*, *gyrB* and *ftsZ* revealed that these strains form a cluster within the Orientalis clade, close to the species *Vibrio tubiashii*. The results of MLSA, the DDH rate and the phenotypic differences with the type strain of *V. tubiashii* supported the differentiation of the Galician isolates as a new subspecies within *V. tubiashii*, for which the name *V. tubiashii* subsp. *europaensis* subsp. nov. is proposed (type strain PP-638<sup>T</sup> = CECT 8136<sup>T</sup> = DSM 7349<sup>T</sup>). The emended description of *V. tubiashii* is included. The pathogenicity assays widen the host range of *V. tubiashii* to add two unreported species, *Venerupis decussata* and *Donax trunculus*, and the described as relatively resistant species *V. philippinarum*.

Hatchery production is currently the only sustainable method of obtaining bivalve spat for the aquaculture industry. Outbreaks of bacterial disease caused by some *Vibrio* species represent the main bottleneck in the production process in bivalve hatcheries. Vibriosis, which is characterized by a sudden large reduction in larval motility and by detached vela and necrotic soft tissue, leads to high mortality rates of more than 90% in 24 – 48 h [14].

The genus *Vibrio* is the largest member of the family *Vibrionaceae* and comprises more than 90 bacterial species (<http://www.bacterio.net/vibrio.html>), classified in 17 clades [20]. Some vibrios have been described as opportunistic pathogens in bivalve aquaculture and may proliferate under culture conditions and kill bivalve larvae [4,8,10,14,16]. The species *Vibrio tubiashii* [6] was recognized almost fifty years ago as a significant pathogen of

bivalve larvae [28,29], causing “bacillary necrosis”. In the last years, Elston et al. (2008) have demonstrated the emerging resurgence of *V. tubiashii* infections in US with dramatic impact on intensive shellfish production.

In this study, we report the taxonomic characterization of six bacterial isolates obtained from bivalve larvae reared in two Galician hatcheries during outbreaks of disease. The pathogenicity, previously demonstrated with strain PP-638 [14], was extended now to the other isolates and with different bivalve species. The phenotypic and genotypic results showed that this group constitutes a new subspecies within the species *V. tubiashii*, in the Orientalis clade, with special significance for shellfish aquaculture.

Microbiological samples were obtained during episodes of mortality that affected bivalve cultures in two hatcheries, and they were processed as in the previous study [14]. All strains included in the present study were isolated from Thiosulphate-Citrate-Bile-Sucrose plates (TCBS, Oxoid). In Hatchery A, isolates PP-654 and PP-660 were obtained from flat oyster larvae and PP-638<sup>T</sup> and PP-635 were isolated from the seawater used in the same culture. Strains PP2-843 and PP2-978 were isolated from small spat ( $\geq 1000 \mu\text{m}$ ) of Manila clam (*Venerupis philippinarum*) reared in Hatchery B. All isolates were cultured in Marine Agar (MA,

**Abbreviations:** DDH, DNA – DNA hybridization; CECT, Colección Española de Cultivos Tipo, Spain; FAME, fatty acid methyl esters; MALDI-TOF MS, matrix-assisted laser desorption/ionization time of flight mass spectrometry; MLSA, multilocus sequence analysis.

\* Corresponding author. Tel.: +34 881816911; fax: +34 881816966.

E-mail address: [ssn.prado@gmail.com](mailto:ssn.prado@gmail.com) (S. Prado).

<sup>1</sup> These authors have contributed equally to this work.

Pronadisa, Lab Conda) for 24 h at 25 °C, and the phenotypic characterization was carried out as previously described [14]. Utilization of 46 sole carbon sources was evaluated in plates containing basal medium and the certain carbon source [3]. Enzymatic activities and additional phenotypic features were determined with API ZYM and API 20E identification kits (bioMérieux) respectively, with the bacteria suspended in saline solution (NaCl 0.85%). Fermentation of carbon sources was tested with API 50CH strips (bioMérieux), with slight modifications [15].

Analysis of FAME was performed according to the Microbial Identifications Systems (MIDI) described by Sasser [18]. The six isolates and the reference type strains *V. tubiashii* CECT 4196<sup>T</sup> and *Vibrio orientalis* CECT 629<sup>T</sup> were cultured previously for 24 h at 25 °C on Trypticase Soy Agar (Pronadisa, Lab Conda) supplemented with 0.5% NaCl (w/v).


The six isolates shared the main phenotypic features of the genus *Vibrio* [13]: motile rods, facultatively anaerobic, Gram-negative, oxidase and catalase positive, susceptible to the vibriostatic agent O/129, able to reduce nitrates to nitrites and to grow on TCBS. These strains showed high phenotypic homogeneity, with variable reactions only for Simmon's citrate (positive for strains PP-660, PP2-843 and PP2-978) and acid production from melibiose (positive for the isolates PP-635 and PP-654).

The fatty acid profiles of the six isolates were very similar. The major fatty acids were summed feature 3 (39.6 ± 1.6, comprising 16:1 $\omega$ 7c and/or 16:1 $\omega$ 6c), 16:0 (17.0 ± 1.9%), summed feature 8 (14.0 ± 2.0%, comprising 18:1 $\omega$ 7c and/or 18:1 $\omega$ 6c), 14:0 (6.1 ± 0.4%) and 16:0-iso (3.4 ± 0.8%) (the complete profile is shown in Table S1).

DNA was extracted and purified from pure isolates by using the Instagene kit (Bio-Rad). A fragment (>1450 bp) of the 16S rRNA gene was amplified and sequenced from the extracted DNA by using bacterial universal primers specific to the 16S rRNA gene (27F and 1510R) [11]. The genes encoding RNA recombinase alpha subunit (*recA*), RNA polymerase alpha subunit (*rpoA*), uridine monophosphate kinase (*pyrH*), gyrase beta subunit (*gyrB*) and a cell-division protein (*ftsZ*) were used for MLSA. The amplification of the housekeeping genes were carried out as previously described [19,26,27,30], and in few cases the annealing temperature was adjusted to provide specific amplification. All the genes were amplified using Ready To Go PCR-beads (Amersham Pharmacia Biotech). The sequences obtained were analyzed using the Lasergene Seqman program (DNASTAR). Comparative analysis was carried out using the EzTaxon-e server [9] and BLAST [1]. Phylogenetic analysis based on the individual and concatenated sequences were performed using MEGA 5.2 software [23], after multiple alignment of data by ClustalW [25]. Distances and clustering with the Neighbour Joining (NJ), Maximum Likelihood (ML) and Maximum Parsimony (MP) algorithms were determined using bootstrap values based on 1000 replications. The sequence data of the closest *Vibrio* species were obtained from GenBank and the online taxonomic scheme for vibrios (<http://www2.ioc.fiocruz.br/vibrio/AVib/Species.html>). Sequences were deposited in the DDBJ/EMBL/GenBank database and the accession numbers are listed in Table S2.

The enterobacterial repetitive intergenic consensus (ERIC-PCR) and repetitive extragenic palindromic (REP-PCR) techniques were used to discriminate bacterial strains. For genotyping, the six isolates and the two reference strains *V. tubiashii* CECT 4196<sup>T</sup> and *V. orientalis* CECT 629<sup>T</sup> were amplified by ERIC-PCR and REP-PCR, as previously described Rodríguez et al. [17].

The DNA fingerprints (Fig. S1) revealed that Galician isolates are different from the type strains of *V. tubiashii* and *V. orientalis*. Two profiles were differentiated in relation to the origin (Hatchery A and B), and the isolates PP-638 and PP2-843 were selected as representative strains of each clone.



**Fig. 1.** Phylogenetic tree based on concatenated sequences of five housekeeping genes *recA*, *rpoA*, *pyrH*, *gyrB* and *ftsZ* and the 16S rDNA obtained by the NJ method. *Photobacterium damsela* was used as outgroup. Sequence data are given in Table S2. Horizontal branch lengths are proportional to evolutionary divergence. Bootstrap ( $\geq 50\%$ ) from 1000 replicates appears next to the corresponding branch. Species belonging to Orientalis clade were included in the grey box. Circles indicate that corresponding nodes were recovered in trees generated with MP method (open circles) and ML method (filled circles). Bar, 0.02 substitutions per nucleotide position.

Phylogenetic analysis based on the 16S rRNA gene sequences using the NJ, MP and ML algorithms (Fig. S2) indicated that strains PP-638 and PP2-843 formed a cluster within the genus *Vibrio*. On the basis of these 16S rDNA sequences, the species closest to strain PP-638<sup>T</sup> were *V. orientalis* CIP 102891<sup>T</sup> (98.46%), *Vibrio mytili* CECT 632<sup>T</sup> (98.26%), *Vibrio owensii* DY05<sup>T</sup> (98.24%), *Vibrio tapetis* CECT 4600<sup>T</sup> (98.23%) and *V. tubiashii* ATCC 19109<sup>T</sup> (98.01%). These percentages are all lower than the threshold value (98.70%) proposed by Stackebrandt and Ebberts [22] for defining new species.

The new cluster was maintained in the individual phylogenetic tree of each housekeeping gene (Fig. S3). However, the bivalve pathogen *V. tubiashii* was the most closely related species according to the phylogenetic analysis and the similarities of all the housekeeping gene sequences: *recA* (95.6 – 95.8%, range of lengths, 817 – 822 bp), *rpoA* (99.1 – 99.2%, 860 – 925 bp), *pyrH* (97.8 – 97.9%; 496 – 560 bp), *gyrB* (96.4%, 1050 – 1109 bp) and *ftsZ* (99.6 – 99.7%, 570 – 575 bp). These values were above the species limits established for some of these genes: 98% for *rpoA*, 94% for *recA* and *pyrH* [26].

The phylogenetic trees based on concatenated sequences (4878 bp) of the 16S rRNA and the five housekeeping genes *ftsZ*, *gyrB*, *pyrH*, *recA* and *rpoA*, determined using NJ, MP and ML algorithms (Fig. 1), confirmed the close relationship between the Galician isolates and *V. tubiashii*. Distance matrices for the MLSA combined sequence dataset (data not shown) provided additional evidences, with similarities of 97.6% with the type strain of *V. tubiashii*, clearly higher than the value proposed by Thompson et al. [24] for delimiting the species of genus *Vibrio* (95%). The similarities with the other members of the Orientalis clade were in the range 90.9 – 93.0%, within the specific concatenated identity between species established by Sawabe et al. [20] for the Orientalis clade (88.8 – 93.7%) using eight gene MLSA.

The DDH experiments were carried out by the hydroxyapatite method in microtitre plates [31]. The experiments were performed by duplicate. Strain PP-638 was hybridized with the isolate PP2-843, and the reference strains *V. tubiashii* CECT 4196<sup>T</sup> and *V. orientalis* CECT 629<sup>T</sup>. The DDH reassociation value within the cluster, between PP-638 and PP2-843, was 87.63%. The degree of

**Table 1**

Features useful in differentiating Galician isolates from *V. tubiashii* and *V. orientalis*. Taxa 1, *V. tubiashii* subsp. *europaensis* (Galician isolates,  $n = 6$ ); 2, *V. tubiashii* subsp. *tubiashii* CECT 4196<sup>T</sup>; 3, *V. orientalis* CECT 629<sup>T</sup>. All data from this study. + positive; – negative.

Features:	1	2	3
Lysine decarboxylase	–	–	+
<b>Growth at/with:</b>			
4 °C	–	–	+
35 °C	–	+	+
8% NaCl	–	–	+
<b>Enzymatic activities (API ZYM)</b>			
Esterase lipase (C8)	+	+	–
Lipase (C14)	+	–	–
Valine arylamidase	–	+	+
Cystine arylamidase	–	+	–
Acid phosphatase	–	–	+
Naphthol-AS-β1-phosphohydrolase	–	+	+
β-galactosidase	–	+	–
<b>Utilization</b>			
d-galactose	+	+	–
Aspartic acid	–	+	+
L-citruline	+	+	–
Myoinositol	–	–	+
Putrescine	–	–	+
Tyrosine	+	+	–
<b>Fermentation</b>			
Arbutine	–	–	+
Salicine	+	–	+
d-turanose	–	+	+

hybridization between PP-638<sup>T</sup> and the reference type strain of *V. tubiashii* CECT 4196<sup>T</sup> was 66.55%, similar to the values recorded between subspecies of *V. tapetis* [2]. The DDH rate with *V. orientalis* CECT 629<sup>T</sup> was 59.95%.

Several phenotypic characters differentiated Galician isolates from the type strains of *V. tubiashii* and *V. orientalis* (Table 1). These included growth at 35 °C, the activities of lipase (C14), valine arylamidase and naphthol-AS-β1-phosphohydrolase, and the fermentation of d-turanose. In addition, the Galician isolates and *V. tubiashii* were also distinguishable in the results of utilization of aspartic acid, fermentation of salicine, and the activities of cystine arylamidase and β-galactosidase (API ZYM).

The MALDI-TOF-MS profile of the strain PP-638 was obtained at CECT, following Maier et al. [12]. The protein analysis confirmed that the closest species were *V. tubiashii* CECT 4196<sup>T</sup> (score = 2.350), in the species-limit value (2.300) [4], and *V. orientalis* CECT 629<sup>T</sup> (score = 1.979).

The FAME profiles obtained (Table S2) demonstrated the similarity between the Galician isolates and *V. tubiashii*. They were useful for distinguishing them from *V. orientalis*, according to the fatty acids iso-C16:0, iso-C17:0 and iso-C15:0, not detected or detected only in trace quantities in *V. orientalis*.

Experimental infections of bivalve larvae were carried out as in a previous study [14]. The pathogenicity of the six Galician isolates was evaluated in four different challenges performed respectively with larvae (10 to 15 days old) of the original hosts, flat oyster and Manila clam, the traditionally reared in Galicia carpet-shell clam *Venerupis decussata*, and the emerging cultured wedge-shell clam *Donax trunculus*. The strains were assayed individually with each bivalve species at a final concentration of 10<sup>5</sup> – 10<sup>6</sup> CFU/ml. The assays were carried out in triplicate and included a negative control without addition of bacteria.

The six isolates caused mortalities of 100% in 48 h in all experimental infections, whereas the survival rates were ≥95% in the negative controls. The pathogens were successfully re-isolated after each challenge as pointed the Koch's postulate.

Our results confirmed the pathogenicity of the Galician isolates, previously demonstrated for strain PP-638 [14], and the

broad action range to kill the larvae of different bivalve species. *V. tubiashii* causes similar effects in bivalve larval cultures, and its re-emergence has recently been reported in different shellfish hatcheries [5]. In this study, we demonstrated the presence of a different subspecies of *V. tubiashii*, which affect the production of bivalve hatcheries of the NW Atlantic coast of Spain. The subspecies resulted pathogenic for *Ostrea edulis*, as reported for *V. tubiashii* [7,28], for previously unreported susceptible species, *V. decussata* and *D. trunculus*, and for *V. philippinarum*, described as relatively resistant to *V. tubiashii* [5].

In summary, the results provide clear evidence that Galician isolates form a cluster closely related to the type strain *V. tubiashii*, but distinguishable in some phenotypic features. MLSA and DDH values demonstrated the separation of this cluster, and following the recommendations of the *ad hoc* committee on reconciliation of approaches to bacterial systematics [21], the new subspecies *V. tubiashii* subsp. *europaensis* subsp. nov. is proposed to include the Galician isolates.

### Emended description of *Vibrio tubiashii*

The characteristics added to those reported in the original description by Hada et al. [6] are as follows.

The major fatty acids are as follows: summed feature 3 (comprises C<sub>16:1</sub> ω7c and/or C<sub>16:1</sub> ω6c), C<sub>16:0</sub>, summed feature 8 (C<sub>18:1</sub> ω7c and/or C<sub>18:1</sub> ω6c), C<sub>14:0</sub>, iso-C<sub>16:0</sub>, C<sub>12:0</sub>, summed feature 2 (C<sub>14:0</sub> 3-OH and/or iso-C<sub>16:1</sub> I), C<sub>12:0</sub> 3-OH, iso-C<sub>17:0</sub>, iso-C<sub>13:0</sub> and iso-C<sub>15:0</sub>. In enzymatic activity tests, *V. tubiashii* is positive for alkaline phosphatase, esterase lipase (C8), leucine arylamidase and N-acetyl-β-glucosaminidase, and negative for esterase (C4), trypsin, α-chymotrypsin, acid phosphatase, α-galactosidase, β-glucuronidase, α-glucosidase, β-glucosidase, α-mannosidase and α-fucosidase. *V. tubiashii* ferments ribose, but not d-arabinose, L-xylose, methyl-β-d-xylopyranoside, L-sorbose, methyl-α-d-mannopyranoside, amygdalin, inulin, d-melezitose, xylitol, d-lyxose, d-tagatose, d-fucose, L-fucose, L-arabitol or potassium 5-keto-gluconate. *V. tubiashii* utilizes d-ribose, L-serine, β-hydroxybutyric acid, lactic acid, succinic acid, propionic acid, malic acid, glycerol, L-threonine, fumaric acid, sodium acetate, maltose, d-fructose, pyruvate, L-histidine, N-acetylglucosamine, d-mannitol, ornithine, L-citruline, L-arginine and citric acid as sole carbon sources. None of the strains can utilize lactose, d-saccharic acid, amygdalin, arabinose, myoinositol, trans aconitic acid, L-leucine, lysine or L-rhamnose as sole carbon sources.

*V. tubiashii* subsp. *tubiashii* subsp. nov. Hada et al. [6].

(Type strain = ATCC 19109<sup>T</sup>)

*V. tubiashii* subsp. *tubiashii* characteristics are those stated in the emended description of *V. tubiashii*.

*V. tubiashii* subsp. *europaensis* subsp. nov. [eu.ro.pa.en'sis. L. adj. of/from Europa].

Characteristics of this subspecies are as follows.

Colonies are round with smooth margins, greenish beige in colour and non-swarming on MA plates. Isolates are not able to grow at 35 °C. None strain utilizes aspartic acid. All strains ferments salicine but no d-turanose. In enzymatic activity tests, the strains are positive for lipase (C14) and they are negative for valine arylamidase, cystine arylamidase, naphthol-AS-β1-phosphohydrolase and β-galactosidase.

The type strain PP-638<sup>T</sup> (=CECT 8136<sup>T</sup> = DSM 7349<sup>T</sup>) was isolated during episodes of mortality affecting larval cultures of flat oyster in a shellfish hatchery in Galicia (NW Spain).

### Acknowledgements

The authors thank Soledad Nuñez (Instituto de Acuicultura, Universidad de Santiago de Compostela) for determining the FAME

profiles. This study was financially supported by BAPOHATCH, Secretaría de Estado de I+D+i, Ministerio de Economía y Competitividad (AGL2011–29765).

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.syapm.2014.11.005>.

## References

- [1] Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., Lipman, D.J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein databases search programs. *Nucleic Acids Res.* 25, 3389–3402.
- [2] Balboa, S., Romalde, J.L. (2013) Multilocus sequence analysis of *Vibrio tapetis*, the causative agent of Brown Ring Disease: description of *Vibrio tapetis* subsp. *britannicus* subsp. nov. *Syst. Appl. Microbiol.* 36, 183–187.
- [3] Baumann, P., Baumann, L. (1981) The marine Gram-negative eubacteria: genera *Photobacterium*, *Beneckea*, *Alteromonas*, *Pseudomonas* and *Alcaligenes*. In: Starr, M.P., Stolp, H., Truper, H.G., Balows, A., Schlegel, H.G. (Eds.), *The prokaryotes*, Springer-Verlag, Berlin, pp. 1302–1330.
- [4] Beaz-Hidalgo, R., Diéguez, A.L., Cleenwerck, I., Balboa, S., Doce, A., de Vos, P., Romalde, J.L. (2010) *Vibrio celticus* sp. nov., a new *Vibrio* species belonging to the Splendidus clade with pathogenic potential for clams. *Syst. Appl. Microbiol.* 3, 215–222.
- [5] Elston, R.A., Hasegawa, H., Humphrey, K.L., Polyak, I.K., Häse, C.C. (2008) Re-emergence of *Vibrio tubiashii* in bivalve shellfish aquaculture: severity, environmental drivers, geographic extent and management. *Dis. Aquat. Org.* 82, 119–134.
- [6] Hada, H.S., West, P.A., Lee, J.V., Stemmler, J., Colwell, R.R. (1984) *Vibrio tubiashii* sp. nov., a pathogen of bivalve mollusks. *Int. J. Syst. Bacteriol.* 34, 1–4.
- [7] Jeffries, V.E. (1982) Three *Vibrio* strains pathogenic to larvae of *Crassostrea gigas* and *Ostrea edulis*. *Aquaculture* 29, 201–226.
- [8] Kesarcodi-Watson, A., Kaspar, H., Lategan, M.J., Gibson, L. (2009) Two pathogens of Greenshell mussel larvae, *Perna canaliculus*: *Vibrio splendidus* and *V. coralliilyticus/neptunius*-like isolate. *J. Fish Dis.* 32, 499–507.
- [9] Kim, O.S., Cho, Y.J., Lee, K., Yoon, S.H., Kim, M., Na, H., Park, S.C., Jeon, Y.S., Lee, J.H., Yi, H., Won, S., Chun, J. (2012) Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *Int. J. Syst. Evol. Microbiol.* 62, 716–721.
- [10] Lambert, C., Nicolas, J.L., Cilia, V., Corre, S. (1998) *Vibrio pectenica* sp. nov., a pathogen of scallop (*Pecten maximus*) larvae. *Int. J. Syst. Bacteriol.* 48, 481–487.
- [11] Lane, D.J. (1991) 16S/23S rRNA sequencing. In: Stackebrandt, E., Goodfellow, M. (Eds.), *Nucleic Acid Techniques in Bacterial Systematics*, Wiley, London, pp. 115–175.
- [12] Maier, T., Klepel, S., Renner, U., Kostrzewa, M. (2006) Fast and reliable MALDI-TOF MS based microorganism identification. *Nat. Methods* 3, <http://dx.doi.org/10.1038/nmeth870>.
- [13] Nogueroles, I., Blanch, A.R. (2008) Identification of *Vibrio* spp. with a set of dichotomous keys. *J. Appl. Microbiol.* 105, 175–185.
- [14] Prado, S., Romalde, J.L., Montes, J., Barja, J.L. (2005) Pathogenic bacteria isolated from disease outbreaks in shellfish hatcheries. First description of *Vibrio neptunius* as an oyster pathogen. *Dis. Aquat. Org.* 67, 209–215.
- [15] Prado, S. 2006 *Microbiota Asociada a Criaderos de Moluscos: Patogénesis y Probiosis* (Ph.D. thesis), Department of Microbiology and Parasitology, University of Santiago de Compostela, Santiago de Compostela, Spain.
- [16] Prado, S., Dubert, J., Romalde, J.L., Toranzo, A.E., Barja, J.L. (2014) *Vibrio ostreicida* sp. nov., a new pathogen of bivalve larvae. *Int. J. Syst. Evol. Microbiol.* 64, 1641–1646.
- [17] Rodríguez, J.M., López-Romalde, S., Beaz, R., Alonso, M.C., Castro, D., Romalde, J.L. (2006) Molecular fingerprinting of *Vibrio tapetis* strains using three PCR-based methods: ERIC-PCR, REP-PCR and RAPD. *Dis. Aquat. Org.* 69, 175–183.
- [18] Sasser, M. 1990 Identification of Bacteria by Gas Chromatography of Cellular Fatty Acids, Microbial ID, Inc, Newark, Delaware.
- [19] Sawabe, T., Kita-Tsukamoto, K., Thompson, F.L. (2007) Inferring the evolutionary history of vibrios by means of Multilocus Sequence Analysis. *J. Bacteriol.* 21, 7932–7936.
- [20] Sawabe, T., Ogura, Y., Matsumura, Y., Feng, G., Amin, A.R., Mino, S., Nakagawa, S., Sawabe, T., Kumar, R., Fukui, Y., Satomi, M., Matsushima, R., Thompson, F.L., Gómez-Gil, B., Christen, R., Maruyama, F., Kurokawa, K., Hayashi, T. (2013) Updating the *Vibrio* clades defined by multilocus sequence phylogeny: proposal of eight new clades, and the description of *Vibrio tritonius* sp. nov. *Front. Microbiol.* 4, 414.
- [21] Stackebrandt, E., Frederiksen, W., Garrity, G.M., Grimont, P.A.D., Kämpfer, P., Maiden, M.C.J., Nesme, X., Roselló-Mora, R., Swings, J., Trüper, H.G., Vauterin, L., Ward, A.C., Whitman, W.B. (2002) Report of the *ad hoc* committee for the reevaluation of the species definition in bacteriology. *Int. J. Syst. Evol. Microbiol.* 52, 1043–1047.
- [22] Stackebrandt, E., Ebers, J. (2006) Taxonomic parameters revisited: tarnished gold standards. *Microbiol. Today* 33, 152–155.
- [23] Tamura, K., Peterson, D., Stecher, G., Nei, M., Kumar, S. (2011) MEGA5: molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28, 2731–2739.
- [24] Thompson, C.C., Vicente, A.C.P., Souza, R.C., Vasconcelos, A.T.R., Vesth, T., Alves, N., Jr., Ussery, D.W., Iida, T., Thompson, F.L. (2009) Genomic taxonomy of vibrios. *BCM Evol. Biol.* 9, 258.
- [25] Thompson, J.D., Higgins, D.G., Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22, 4673–4680.
- [26] Thompson, F.L., Gevers, D., Thompson, C.C., Dawyndt, P., Naser, S., Hoste, B., Munn, C.B., Swings, J. (2005) Phylogeny and molecular identification of vibrios on the basis of multilocus sequence analysis. *Appl. Environ. Microbiol.* 71, 5107–5115.
- [27] Thompson, F.L., Gómez-Gil, B., Vasconcelos, A.T., Sawabe, T. (2007) Multilocus sequence analysis reveals that *Vibrio harveyi* and *V. campbellii* are distinct species. *Appl. Environ. Microbiol.* 73, 4279–4285.
- [28] Tubiash, H.S., Chanley, P.E., Leifson, E. (1965) Bacillary necrosis, a disease of larval and juvenile bivalve mollusks. I. Etiology and epizootiology. *J. Bacteriol.* 90, 1036–1044.
- [29] Tubiash, H.S., Colwell, R.R., Sakazaki, R. (1970) Marine vibrios associated with bacillary necrosis, a disease of larval and juvenile bivalve mollusks. *J. Bacteriol.* 103, 272–273.
- [30] Yamamoto, S., Harayama, S. (1995) PCR amplification and direct sequencing of *gyrB* genes with universal primers and their application to the detection and taxonomic analysis of *Pseudomonas putida* strains. *Appl. Environ. Microbiol.* 61, 1104–1109.
- [31] Ziemke, F., Höfle, M.G., Lalucat, J., Roselló-Mora, R. (1998) Reclassification of *Shewanella putrefaciens* Owen's genomic group II as *Shewanella baltica* sp. nov. *Int. J. Syst. Bacteriol.* 48, 179–186.

